



---

## **VESSEL SPILL RESPONSE TECHNOLOGIES**

---



---

## **GULF OF THE FARALLONES AND CORDELL BANK NATIONAL MARINE SANCTUARIES**

---

*Report of a Joint Working Group of the Gulf of the Farallones and Cordell Bank  
National Marine Sanctuaries Advisory Councils*

**June 2012**

*We would like to thank:*

PRBO Conservation Science for supporting the working group's meetings and the institutions, agencies, and organizations that supported their staff and representatives in participating.



prbo

PRBO Conservation Science

## **Working Group Members**

---

Yvonne Addassi, Office of Spill Prevention and Response, CA Dept. of Fish and Game  
Sarah Allen, Ocean Stewardship Program, National Park Service  
Richard Charter, Ocean Foundation; GFNMS Advisory Council  
Barbara Emley, San Francisco Community Fishing Association; GFNMS Advisory Council  
Ellen Faurot-Daniels, Office of Spill Prevention and Response, CA Dept. of Fish and Game  
Jaime Jahncke (Chair), PRBO Conservation Science; GFNMS and CBNMS Adv. Councils  
Gerry McChesney, Farallon National Wildlife Refuge, US Fish and Wildlife Service  
Renee McKinnon, US Coast Guard - Sector San Francisco (June 2011- February 2012)  
James Nunez, US Coast Guard - Sector San Francisco (March 2012 - June 2012)  
Patrick Rutten (Chair), NOAA Restoration Center; GFNMS Advisory Council  
Deb Self, San Francisco Baykeeper (June 2011- February 2012)  
Jordan Stout, Office of Response and Restoration, NOAA  
Bob Wilson, Farallones Marine Sanctuary Association; Marine Mammal Center;  
GFNMS Advisory Council  
Ian Wren, San Francisco Baykeeper (March 2012 - June 2012)

## **Invited Agency Observers**

---

Leslie Abramson, Gulf of the Farallones National Marine Sanctuary, NOAA  
Maria Brown, Superintendent, Gulf of the Farallones National Marine Sanctuary, NOAA  
Joseph Dillon, National Marine Fisheries Service, NOAA  
Steve Gittings, Office of National Marine Sanctuaries, NOAA  
Doug Helton, Office of Response and Restoration, NOAA  
Dan Howard, Superintendent, Cordell Bank National Marine Sanctuary, NOAA  
John Hunt, Monterey Bay National Marine Sanctuary, NOAA  
Scott Kathey, Monterey Bay National Marine Sanctuary, NOAA  
Dave Lott, Office of National Marine Sanctuaries, West Coast Region, NOAA  
Bill Robberson, Environmental Protection Agency  
Jan Roletto, Gulf of the Farallones National Marine Sanctuary, NOAA  
GP Schmahl, Flower Garden Banks National Marine Sanctuary, NOAA  
Alice Stratton, Stellwagen Bank National Marine Sanctuary, NOAA  
Lisa Symons, Office of National Marine Sanctuaries, NOAA

## **Supporting Staff**

---

Irina Kogan, Gulf of the Farallones National Marine Sanctuary, NOAA  
Michael Carver, Cordell Bank National Marine Sanctuary, NOAA  
Meredith Elliott, PRBO Conservation Science

## Table of Contents

---

<b>EXECUTIVE SUMMARY</b>	<b>Page 1</b>
<b>INTRODUCTION</b>	<b>4</b>
Purpose and Process	4
<b>BACKGROUND</b>	<b>5</b>
Sanctuary Role in Spill Response	5
On-Water Response Technologies	5
Net Environmental Benefits Analysis (NEBA)	8
Toxicity of Oil and Oil/Dispersant Mixtures	11
Oceanography of North-Central California	16
Biological Resources in the Sanctuaries	17
<b>RECOMMENDATIONS</b>	<b>21</b>
Science and Research Recommendations	21
Education and Outreach Recommendations	22
Policy and Management Recommendations	22
Specific Sanctuary Recommendations	23
<b>APPENDICES</b>	<b>24</b>
Appendix I: Meeting Agendas	24
Appendix II: Working Group Member Biographies	26
Appendix III: Invited Speaker Biographies	30
Appendix IV: List of Marine Species of GFNMS and CBNMS	34
Appendix V: Sensitive Species Matrix	78

## Executive Summary

---

In June 2011 the Superintendents of the Gulf of the Farallones (GF) and Cordell Banks (CB) National Marine Sanctuaries (NMS) established a Vessel Spills Working Group (VSWG) to provide an analysis and advisory report on the use of oil dispersants within the GF and CB NMS. The formation of the VSWG was the result of a recommendation in the GFNMS 2008 Management Plan. The objective of the VSWG is to provide a set of recommendations to the Sanctuary Advisory Councils (SAC) to consider and transmit to the Superintendents for their consideration. This process was completed in May 2012 and report presented to the SACs on June 7, 2012.

The VSWG had invited technical members who discussed inter-agency coordination and response, dispersant decision protocols, oil spill trajectory models, and response technologies. The VSWG decided to focus more on oil spill response technologies and specifically the use of dispersants. In order to fully understand the complexity and dynamics of the fate of oil and oil dispersants and the potential impacts of dispersed oil on the resources within the Sanctuaries, the VSWG has conducted a series of meetings with presentations from identified regional experts in the areas of toxicology, oceanography and the biological resources of the Sanctuaries. The effect of oil and dispersed oil on human health was not a topic discussed with the VSWG. However, a general discussion on this topic is included at the end of the background section based on a group consensus that the topic merits consideration.

The following are key points from this report:

Sanctuaries **are not** within a dispersant pre-approval zone.

### **Sanctuary Role in Spill Response**

The Office of National Marine Sanctuaries has a consultative role in these decisions. The Sanctuary Superintendent does not make the final decision.

### **On-water Response Technologies**

Three on-water response options: mechanical recovery (skimming), *in-situ* burning, chemical dispersants. Mechanical recovery rates are typically less than 20% in sheltered waters and often less than 10% in open-water. *In-situ* burning requires seas less than 2–3 ft. (0.6–0.9 m). Oceanic and regulatory limitations on *in-situ* burning limit its use as a primary oil spill response option in California. Effective chemical dispersion of oil requires surface mixing energy (typically a few knots of wind and a light chop). Dispersant operations encounter rates are 10-100 times greater than skimming or burning. Other than no action, dispersants may be the only response option during rough, open-water conditions. Chemically dispersed oil may adversely impact organisms in the upper water column.

### **Net Environmental Benefits Analysis (NEBA)**

A Net Environmental Benefit Analysis (NEBA) uses a risk matrix to evaluate scenario-based comparisons of different response strategies and their associated environmental tradeoffs. The risk matrix provides a qualitative ranking of population percentage impacted and expected recovery time.

### **Toxicity of Oil and Oil/Dispersant Mixtures**

Use of chemical dispersants does introduce higher total concentrations of petroleum hydrocarbons into the water column than naturally dispersed oil. This higher concentration may have a larger footprint to potentially impact a wider range of species that would not likely have been exposed or affected by the surface oil slick. Nearly all chemicals are toxic at some concentration, so to say that a particular chemical or chemical mixture is “toxic” may not necessarily be true at environmentally-relevant concentrations.

Embryo-larval stages and early juvenile life stages are generally more sensitive to chemicals than are adults of the same species. Water containing dispersed oil droplets and oil that reaches the gills of fish can potentially cause effects. Different organisms and life stages have varying sensitivities. Many California endemic species have been used in toxicity studies involving oil and dispersants (including red abalone, giant kelp, mysids shrimp, Chinook salmon and top smelt). Species of concern found in the Gulf of the Farallones that have not had toxicity test data include black abalone and Dungeness crab.

### **Oceanography of North-Central California**

Transport of surface oil and subsurface (i.e. dispersed) oil may be different based upon wind and current patterns at the time of the spill. Oil dispersed into deeper water will move with midwater currents, while oil at the surface will move with the surface currents as influenced by winds and may move onshore.

During times of upwelling, it is expected dispersed oil will remain in the upper water column, while during times of downwelling, dispersed oil will be driven deeper into the water column where it will experience significant dilution. It is expected some dispersed oil will travel into the nearshore zone during downwelling. Unlike Southern California, there has been no regional current forecast modeling for Northern California.

### **Biological Resources in the Sanctuaries**

Biological resources were evaluated for the potential negative effects from dispersants ranging from the simplest plankton to birds and mammals and we assembled a list of species of interest drawn from the larger list of species that occur in the region (Appendix IV). Ultimately, dispersant-use decisions will be guided by the potential percentage impact on the population and recovery time of a species.

### **Invertebrates**

Most zooplankton populations are not likely to be permanently affected by oil spills and are expected to recover due to their high population numbers and wide distribution. Larval stages of invertebrates and fish are considered susceptible to oil or dispersants in the water column if exposed. For many invertebrates, the adult phase is considered a high priority for protection because of their reproductive capability. It is expected that individual larvae will be lost but population-level effects will be unlikely.

### **Fish**

Adult salmon on their landward migration are less susceptible to dispersed oil exposure due to their generally rapid movement into San Francisco Bay and ability to swim quickly. Juvenile out-migrating salmon are potentially more vulnerable to oil and dispersed oil due to increased residency time in the GF and lower general swim speeds.

Rockfish are found wherever suitable habitat is located in the Sanctuaries. Rockfish do not move widely and are considered more vulnerable to oil spills locally, but are generally found at depths that provide significant dilution for dispersed oil and they would be replaced by natural recruitment of animals from adjacent areas.

Wide ranging species with large populations such as anchovies were not considered to be vulnerable to spills or dispersants at the population level. Research has demonstrated that herring eggs and larvae exposed to undispersed oil in the intertidal zone in SF Bay experienced significant mortality that was accelerated by sunlight.

### **Bird and Mammals**

Indirect effects to birds may include accumulation of toxic components from their food, exposure to secondary chemicals (dispersants), and destruction of habitat or prey resources.

Some sea birds are attracted to surface oil slicks on the water because they look like fish oil slicks. Storm-petrels may be inadvertently attracted to sulfurous crude oil slicks because that particular oil smells like krill (on which they feed) that emit similar compounds.

There is much information on the potential effects of oiling on birds but little information on the effects of dispersants or dispersed oil on feathers or ingestion at environmentally-realistic concentrations.

For mammals, all breeding species are potentially vulnerable to oil spills because of nursing pups/calves that might ingest oil and because most species congregate during feeding. The species most vulnerable to exposure to oil are those that rely on fur for insulation including sea otters and fur seals.

## Introduction

---

### **Purpose of the Working Group**

There is a continuing risk of vessel spills that could impact marine mammals, seabirds, other biota, and cultural resources in and around the Gulf of the Farallones National Marine Sanctuary (GFNMS) and Cordell Bank National Marine Sanctuary (CBNMS). Historically, spills have occurred from transiting or sunken vessels with crude oil, bunker fuel, and/or other hazardous material onboard. These incidents have generally been discrete in time and place with the exception of the SS Jacob Luckenbach, a sunken vessel that was addressed once it was identified as the source of periodic oil releases. There are no oil platforms or other potential sources of repetitive spills in the GFNMS or CBNMS at this time. The 2008 GFNMS Management Plan recommended the creation of a vessel spills working group to aid the Sanctuary in understanding and minimizing spill-related risk to Sanctuary natural resources.

The purpose of the Vessel Spills Working Group on Oil Spill Response Technologies (Working Group or VSWG) was to engage agency responders, government resource trustees, and stakeholders such as conservation NGOs and commercial fishing interests in developing a set of recommendations for the Sanctuary Advisory Councils (SACs) to consider in making their recommendations to the GFNMS and CBNMS regarding the use of response technologies in the respective Sanctuaries.

The recommendations from the VSWG are forwarded to the GFNMS and CBNMS SACs for consideration. The SACs will then develop their recommendations on the use of spill response technologies in the respective Sanctuaries to the GFNMS and CBNMS Superintendents. These recommendations will be considered by the GFNMS and/or CBNMS, as appropriate, during a spill in making its recommendation(s) to the Unified Command and the NOAA representative to the Region IX Regional Response Team (RRT-9).

### **Process of the Working Group**

To accomplish its purpose, the Working Group has met seven times and had one conference call between June 2011 and May 2012. Meeting topics included: Response Technologies 101, Net Environmental Benefits Analysis, Dispersants 101 and decision making, Dispersant Toxicity, Sanctuary Biological resources, and Sanctuary Oceanographic Setting. The Working Group has sought to achieve consensus (and record other positions) in the recommendations for the SAC to the fullest extent possible. The Working Group consists of a body of individuals representing diverse interests and perspectives (Appendix II). In addition to obtaining technical information and expertise to develop recommendations, members have engaged in meaningful dialogue and informed/educated their constituency groups.

## Background

---

### **Sanctuary Role in Spill Response**

Under the authority of the federal Oil Pollution Act of 1990, primary response responsibility for marine spills has been delegated to the U.S. Coast Guard (USCG). Therefore, the USCG is the lead federal agency on all marine spill response and planning activities. During a spill event, the Unified Command (UC) orchestrates all emergency response and cleanup activities and consists of the USCG (federal), the State of California's Office of Spill Prevention & Response (OSPR) if the spill is within or threatens state waters and/or resources, and the Responsible Party (RP). The UC may choose to bring in other agencies and/or private parties to assist during an event, including GFNMS and CBNMS as resource trustees. The UC establishes and oversees an Incident Command (IC) staff that is comprised of individuals with unique expertise from a variety of organizations.

A California Dispersant Plan and Federal On-Scene Coordinator checklist have been developed for determining the feasibility of using dispersants in California and can be found in the Regional Contingency Plan for Region 9 (California, Nevada, and Arizona) (2008 RCP). Dispersants have been "pre-approved" by the Region 9 Response Team (RRT-9) for use outside of Sanctuaries or beyond 3 nautical miles of any landfall (state waters) or Mexico. Pre-approval requires that the USCG follow the dispersant pre-approval checklist and ensure all criteria have been met. If this is the case, further consultation with the RRT-9 is not required. If, however, all criteria are not met, pre-approval is not authorized and RRT-9 approval shall be required. Since California National Marine Sanctuaries are not in the pre-approval zones, any dispersant use request made by the USCG during a spill in the Sanctuaries will require approval by the RRT-9.

Although the Sanctuary Superintendent does not make the final decision on whether dispersants or other applied technologies (e.g. in-situ burning) are used within the Sanctuaries, Sanctuary staff will be expected to provide input into the decision process through participation in the spill response and through the NOAA representative on the RRT-9. ONMS does not have a vote. ONMS does have a consultative role on major decisions such as use of dispersants, in-situ burning, bioremediation, and shoreline clean-up agents.

### **On-water Response Technologies**

Typically three types of offshore response strategies may be deployed or considered for deployment during an oil spill in order to efficiently remove oil from the water's surface and to prevent the migration of oil to sensitive nearshore and shoreline habitats. Implementation of these strategies is based on San Francisco-Bay Delta Area Contingency Plan (SFBD ACP) and the Region 9 Regional Contingency Plan (RCP). ACPs are developed with input from numerous local, state and federal participants, industry,

oil spill response organizations (OSROs) and non-government organizations (NGOs) and are frequently tested during various drills and exercises. ACPs are local in scope, and address sensitive resources, response strategies and general oil spill response concerns within a certain number of coastal counties (there are several ACPs to cover the entire coast of California). The RCP is, in contrast, a single document authored and maintained primarily by the federal and state members of RRT-9, which includes federal and state natural resource trustee agencies. The RCP addresses response actions that are statewide in nature, and therefore includes plans such as the one for dispersants that do not vary within a region. On-water response operations addressed by ACPs and the RCP generally include mechanical removal (on-water skimming), *in-situ* burning, and the application of chemical dispersants. All three include their own “window of opportunity” and unique set of operational constraints and ecological considerations. Each strategy is implemented with specific operational requirements to minimize impacts to sensitive resources to the greatest extent possible.

“Windows of opportunity” are the timeframes during a spill event when each response method works the best. Variables that influence the window of opportunity include, but are not limited to, the type of material spilled, location, oceanographic and weather conditions, product weathering, emulsion rates, and the different environments, species, and ecosystems that may be impacted. When response strategies are used within these windows, they are more effective. Selecting response options (including natural recovery) involves considering tradeoffs among predicted effectiveness, potential environmental impact, appropriateness for habitat, and timing.

The following discussion focuses only on the three primary on-water response strategies, and is drawn from NOAA’s *Characteristics of Response Strategies: A Guide for Spill Response Planning in Marine Environments* job aid (available at [www.response.restoration.noaa.gov](http://www.response.restoration.noaa.gov)) and other sources.

### **Mechanical Removal (or skimming)**

On-water skimming operations in the Gulf of the Farallones would involve the use of slow-moving, relatively large vessels in conjunction with floating containment boom and surface skimming pumps to mechanically collect, skim and remove oil from the water’s surface.

There are numerous types of skimming devices, including: brush, disc, drum, belt, rope mop, sorbent belt, suction, and weir skimmers. They are placed at the oil/water interface to recover, or skim oil from the water’s surface and may be operated from shore, be mounted on vessels, or be completely self-propelled. Because large amounts of water are often collected with the oil, efficient operations require that floating oil be concentrated at the skimmer head using containment boom. Adequate storage of recovered oil/water mixtures must be available, along with suitable transfer capability.

Mechanical removal is most successful in quiet, protected conditions. Recovery rates for

skimming operations can vary but seldom exceed 20% in the most sheltered waters and are often less than 10% in open-water conditions. This is largely because skimming is boom-dependent, requiring very slow speeds (often under one knot) and thus have low encounter rates. It can generate large volumes of oily water waste and even with experienced operators, oil will begin to escape from containment boom in seas greater than 2-3 feet. Skimming requires very slow speeds and constant monitoring to be effective, while the associated ecological impacts are typically expected to be minimal.

### **In-situ Burning**

*In-situ* burn operations in the Gulf of the Farallones and Cordell Bank region would involve the use of slow-moving vessels and fire retardant containment boom to be effective. In addition, these operations utilize numerous spotter and air quality monitoring support craft to minimize potential impacts to human health and the environment.

*In-situ* burning has been extensively researched, tested, and utilized in response to oil spills and is believed to be one of the most efficient ways of removing surface oil. Even so, like skimming operations, *in-situ* burning is a boom-dependent operation, and thus susceptible to the same types of failures when seas exceed 2–3 ft (0.6–0.9 m) in height. Burning would need to be performed early in the spill event when the oil is relatively fresh and can be kept thick enough (at least 1-2 mm thick) to sustain the burn. A pre-approval for *in-situ* burning is in place for marine waters further than 35 nm from the California coast. Closer to shore, due to concerns about air quality, RRT approval is required for use of *in-situ* burning. Each Air District along the California coast has also developed Quick-Approval zones that can be used if winds are blowing parallel to or offshore, and these can factor into the RRT-9 decision about whether an *in-situ* burn very close to shore or on land will be safe to conduct. Thus, oceanic and regulatory limitations on *in-situ* burning limit its use as a primary oil spill response option in California. From an equipment perspective, California does not currently have any of the necessary and specialized fire boom available; the nearest west coast supply of fire boom is in Washington State, with a minimum 24-hr delivery time to California.

As with skimming operations, ecological impacts of vessel and boom operations during an *in-situ* burn would be expected to be minimal because operations are conducted under slow speeds and constant monitoring by numerous vessels. Though there is a possibility that some marine species might become entrapped within a boomed area, proper wildlife monitoring should minimize such potential impacts. Unlike the Gulf of Mexico, where numerous sea turtles rear starting at the hatchling life-stage, the chance of encountering juvenile sea turtles is discountable in the GFNMS and CBNMS. The probability of encountering adult sea turtles is relatively rare, but the required monitoring is expected to prevent impacting them in an *in-situ* burn. However, the possible effects of large volumes of smoke on wildlife and human health are not well known and the toxicological impacts from burn residues have not been evaluated. On water, burn residues may sink.

### **Dispersants**

The surface application of chemical dispersants in the Gulf of the Farallones and Cordell Bank region would likely involve one or more vessels with spray arms and/or helicopters or fixed-wing aircraft. Applications would be guided by spotter aircraft to ensure dispersants are applied to the thickest and freshest areas of oil and to avoid individual marine mammals and concentrations of other wildlife to the greatest extent possible. Trained teams would also be deployed by boat and aircraft to monitor both the effectiveness in dispersing oil into the water column and to measure dispersed oil concentrations (at various upper water depths inside and outside the dispersant area of operations).

The dispersants in use today are relatively effective at dispersing oil into the water column, less toxic than earlier formulations and typically less toxic than the oils they are used to treat. Dispersants reduce the oil/water interfacial tension, making it easier for waves to break up oil into very small droplets, often less than 50-70 micrometers ( $\mu\text{m}$ ); thus enhancing their biodegradation potential. They also prevent dispersed particles from re-coalescing and forming bigger, more buoyant droplets that will float to the surface, re-creating sheens or slicks. To accomplish this, effective chemical dispersion requires a threshold amount of surface mixing energy (typically a few knots of wind and a light chop).

As with in-situ burning, dispersant operations would need to be performed early in the spill event when the oil is relatively fresh, as effectiveness diminishes as the oil spreads and weathers. Dispersant applications are typically only possible during the first few days, at most, before an oil slick moves, spreads and weathers to the point that application would not be effective. Even so, dispersant operations have much higher encounter rates (10-100 times) than skimming or burning. And given the sea state constraints of the other strategies, dispersants may be the only response option during rough, open-water conditions other than the no-action alternative.

Oceanographic conditions, currents, upwelling, and downwelling will influence the spread of the dispersed oil. Ecological impacts from these dispersant operations must be carefully evaluated. Dispersant use within Sanctuaries would require incident-specific RRT-9 approval before use. Until sufficiently diluted, chemically dispersed oil may adversely impact organisms in the upper water column. At the time of this writing, the State of California has determined that such operations should only be considered in waters deeper than 60 ft (approximately 20m) and when the impact of floating oil is determined to be greater than that of dispersed oil on the water-column. Consideration is typically given to avoid directly spraying any wildlife, especially birds or fur-bearing marine mammals.

## Net Environmental Benefits Analysis (NEBA)

In mounting an effective oil spill response, the USCG works with other agencies to direct efforts protecting public health, welfare, and the environment. Once oil is spilled to the ocean there will inevitably be impacts to the environment, no matter what response strategy is employed. Furthermore, response strategies themselves can cause impacts, so understanding the net environmental benefit of different response strategies can be critical to minimizing overall impacts of a spill event (i.e., from oil and response activities) and allow for quicker environmental recovery.

A formalized Net Environmental Benefit Analysis (NEBA) uses a risk matrix to evaluate scenario-based comparisons of different response strategies and their associated environmental tradeoffs. The risk matrix provides a qualitative ranking of impacts to affected resource populations or communities based on magnitude of concern. As in figure 1 (below), population percentage impacted and expected duration of recovery are graphed. In a format that can easily be compared, spill response options including: 1) no action, 2) mechanical cleanup, 3) *in situ* burning, and 4) dispersant use, can be scored and compared. Conducting such a thorough analysis during an actual spill emergency is exceedingly challenging, so many NEBAs have been completed along the California coast as part of the ACP and RCP planning processes and explicitly for the development of the Dispersant Use Plan.

**Figure:** Simplified example of NEBA risk matrix from the USCG’s consensus-Ecological Risk Assessment guidebook.

		RECOVERY PERIOD			
		> 7 years (SLOW) (1)	3 to 7 years (2)	1 to 3 years (3)	< 1 year (RAPID) (4)
% of RESOURCE	> 60% (LARGE) (A)	1A	2A	3A	4A
	40 - 60% (B)	1B	2B	3B	4B
	20 - 40% (C)	1C	2C	3C	4C
	5 - 20% (D)	1D	2D	3D	4D
	0 - 5% (SMALL) (E)	1E	2E	3E	4E

**Legend:** Cells shaded dark gray represent a high level of concern, cells shaded gray represent a moderate level of concern, and cells not shaded represent a limited level of concern.

In the NEBA process, the benefits and risks of each cleanup option are evaluated separately and then compared. However, an effective spill response may use a combination of several available response options. Depending upon the oceanographic conditions, spill location, type of oil spilled the use of dispersants may be considered in conjunction with mechanical cleanup equipment and other response strategies.

The NEBA is one way to look at and understand differing strategies and their associated environmental trade-offs, singly or in combination. The outcomes of such discussions may then be used as “pre-loaded” information before a real spill event occurs so that time-critical decisions can be made more efficiently. They can be a very helpful way of quickly evaluating many of the spill-specific environmental trade-offs associated with the response strategies under consideration. Many of the NEBA workshops in California to date have focused on RRT-9 designated dispersant Pre-approval Areas. If dispersants are being considered in such areas, then responders will benefit from such pre-loaded information.

The Vessel Spills Workgroup meetings have discussed many of the elements commonly part of a focused NEBA process. Generally, evaluating oil spill response options include assessing environmental tradeoffs because each approach can cause impacts and because no single response approach is likely to protect all resources perfectly. As previously discussed, both mechanical cleanup (skimming) and *in situ* burning are relatively slow-moving, boom-dependent operations, and their success in offshore waters may be severely limited by sea conditions and distance offshore. At times, such response techniques may not significantly reduce the risk of spilled oil contacting biological resources at the sea surface or in coastal (*e.g.*, intertidal) regions.

Shoreline cleanup methods may not be available or appropriate for use in some remote or sensitive coastal habitats (*e.g.*, rocky intertidal, marshes, wetlands). Inappropriate use may pose a greater risk to these sensitive habitats and dependent species than the oil itself. In such cases the best option may be to keep the oil from ever reaching such sensitive areas or reducing the amount of oil that reaches those areas.

When used in an appropriate and timely manner, dispersants can remove a significant amount of oil from the water’s surface. Appropriate and timely application includes a number of decision factors.

- While dispersants may measurably reduce the risk of oil to surface and shoreline resources, there may be a short-term increase in impacts to the plankton community in the upper water column.
- Rapid decisions on dispersant use are essential as they must be applied quickly, before the oil significantly weathers, to be effective.
- Oil dispersed into the upper water column will quickly dilute to levels where acute toxic effects are much less likely.
- Few acute toxic effects have been reported for crude oil dispersed into the upper 10 m of well-mixed water (Mearns pers. comm.) although available information is limited.
- Dispersants are not appropriate for use on diesel and gasoline based on volatility and toxicity concerns of the spilled product, and are not approved for dispersant use in California.

Available field data and models indicate that concentrations of dispersed oil within the upper water column would be expected to decline below 1 part per million (ppm) at 10 m depth (Mearns pers. comm..) within hours as the cloud of micro-droplets diffuses in 3-dimensional space and begins to degrade. Within a matter of days, dispersion and biodegradation processes can remove much of the plume of oil droplets from the upper water column, and/or reduce concentrations to non-detectable levels.

In contrast, undispersed and unrecovered oil left on the water's surface in the open ocean will break up into smaller and smaller patches of weathered oil but may still persist at the surface for weeks to months, where it may continue to impact pelagic birds, mammals and perhaps sea turtles. If the oil moves toward shore, it can strand in sensitive coastal habitats (especially intertidal areas) and potentially pose a persistent threat, on a timescale of months to years, to those sensitive coastal habitats and their dependent species and communities. There may be circumstances when it would be acceptable for undispersed oil to come ashore for clean-up (e.g. sandy beach as compared to rocky intertidal) but this needs to be evaluated on a case-by-case basis.

Other NEBA-relevant points were also discussed during other VSWG meetings and will be covered in other parts of this workgroup report.

NEBA or NEBA-like processes can be very helpful in spill planning and pre-loading information to support time-critical USCG response decisions. The Gulf of the Farallones National Marine Sanctuary (GFNMS) is a unique ecosystem which provides a broad range of ecological benefits to a variety of plants, invertebrates, fish, birds, sea turtles and mammals. While the existing NEBAs for the areas within the Sanctuaries provide adequate and sufficient resource detail to make a real time assessment on the potential impacts of using dispersants within the GFNMS, current Sanctuary staff were not involved in developing those NEBAs. As NEBA information can be helpful for Sanctuary management staff during a spill event and will help make USCG operational decisions more robust and protective of the environment in the event of a large spill. It is therefore recommended that Sanctuary staff consider performing a small series of scenario-based NEBA discussions with USCG and subject matter experts to identify environmental decision drivers, existing data gaps and developing additional quick-turnaround inputs to support USCG time-critical response decisions.

### **Toxicity of Oil and Oil/Dispersant Mixtures**

Determining whether or not to support the decision to apply chemical dispersants to maritime oil spills is often one of the most controversial and consequential decisions facing resource managers and spill responders. Their input should therefore be based on objective, science-based information and should weigh the environmental and other trade-offs associated with dispersants and the use of alternative response strategies that may be available.

### **Dispersant Formulations**

How dispersants work, their strengths, limitations and environmental considerations are described elsewhere in this report. Early use of chemicals to treat oil spills was both effective and very toxic. Since that time, chemicals have been specifically formulated to disperse surface oil slicks with a higher level of efficacy while lowering the level of toxicity relative to the targeted oil. The working group discussed a range of variables related to toxicity, including genotoxic and mutagenic properties, and more detail on this discussion is provided below.

When considering the toxicological implications of a dispersant, it is important to remember that a cloud of dispersed particulates/micro-droplets will diffuse in three dimensions. This dispersed oil cloud is expected to dilute to lower concentrations in the water column and would thus not be much influenced by winds that could push the surface oil towards shore. Also, such micro-droplets are more readily biodegraded than naturally dispersed oils larger droplets, by virtue of their high surface to volume ratio. That being said, the use of chemical dispersants does introduce higher total concentrations of petroleum hydrocarbons into the water column than naturally dispersed oil. This higher concentration enters more rapidly into a larger volumetric footprint to potentially impact a greater number of organisms that may not have been affected by or exposed to the undispersed oil slick. The working group discussed the basis of these inputs and how they would be used in a formalized NEBA process.

### **General Toxicology Concepts**

While there is broad agreement and strong data to support the known toxic and mutagenic properties of crude oil and related petroleum products, the range of chemical components represented in various dispersants has long complicated any discussion about the fates and effects of dispersant application in marine ecosystems. These differences of interpretation also emerged during technical presentations to, and deliberations among, the Vessel Spills Working Group. The time and scope of the workgroup process did not allow for the consideration of potential human health impacts of petroleum products and dispersants. However, human health issues were discussed broadly and the group felt these topics should be discussed more thoroughly at another time.

Nearly all chemicals are toxic at some concentration, so to say that a particular chemical or chemical mixture is “toxic” may not necessarily be true at environmentally-relevant concentrations and may therefore skew objective environmental trade-off discussions. Furthermore, both oil and dispersants are complex mixtures of many different chemical components, so a holistic view of a given oil or dispersants real-world toxicity to any one of a huge variety of organisms and habitats can quickly become complicated. Embryo-larval stages and early juvenile life stages are generally more sensitive to chemicals than are adults of the same species.

During the meeting Dr. Tjeerdema indicated that much toxicity testing of dispersants has occurred in CA since 1987 and that interpreting the body of available dispersant/oil toxicity data can be difficult. First, caution should be used to rely primarily on peer-reviewed documents authored by established experts in the fields of toxicology and oil spills since less rigorous, web-based sources of information may be of varying quality and/or may mislead readers by providing good information out of context. However, even peer-reviewed toxicity data can be confusing to many for a variety of reasons including:

- Varying oil/water, dispersant/water, oil/dispersant/water mixture preparations
- How concentrations are reported (e.g. THC, TPH, WAF, CEWAF, etc.)
- How effects are measured (e.g. lethality, sub-lethal, gene expression, metabolic changes, etc.)

Toxicity results are often reported in LC50s (the concentration of pollutant in water that causes death to 50% of the test organisms), EC50s (the concentration of pollutant in water that causes some sub-lethal effect to 50% of the test organisms), NOAEL (No Observable Adverse Effect Level), LOAEL (Lowest Observed Adverse Effect Level). Even with understandable toxicity results, applying them to a real world situation also requires an understanding of reasonable routes of exposure and realistic environmental concentrations and durations because risk to a given organism is a function of chemical concentration in the water, an actual exposure to said concentrations and duration of said exposures. The working group discussed these topics as well as acute and chronic exposures, likely environmental concentrations and exposures over space and time.

The Environmental Protection Agency (EPA) is charged with listing dispersants on the National Contingency Plan (NCP) Product Schedule based on both dispersant effectiveness and toxicity data provided by the manufacturers. Toxicity testing currently associated with listing of a product on the NCP Product Schedule focuses on acute toxicity within 48 and 96 hours after dispersant application. Specifically, this type of constant laboratory exposure regime typically overestimates acute toxicity probability as the concentrations during a spill are not constant over these periods of time and rarely models longer term and/or multigenerational exposures. Longer-term studies may help to better capture the impacts that occur over longer time periods, such as neurotoxicity, cardiovascular toxicity, organ damage, infertility, genetic damage and lesions and tumors.

#### **Route of Exposure (Particulates versus Dissolved)**


While undispersed oil generally poses the greatest threat to shorelines and surface-dwelling organisms, dispersed oil may also threaten water column organisms. Most oils float and though surface slicks will naturally break up over time into floating streamers, patches and then tarballs; even without dispersants some portion will also become naturally dispersed into the water column in the form of small particles or droplets and will also dissolve. When chemically dispersed, a floating oil will enter the water column

as a more concentrated cloud of very small particles or micro-droplets and also as dissolved constituents. Water containing dispersed oil droplets and oil that reaches the gills of fish can potentially cause effects. Interestingly, research shows that the dissolved fraction of oil is similar regardless of dispersant use even though particulate/droplet load is greatly increased with dispersants. This becomes important when considering real-world routes of exposure because particulate-feeding organisms (e.g. zooplankton, small nekton) will have increased exposure to chemically-dispersed oils than naturally dispersed oils whereas non-ingesters (e.g. phytoplankton, eggs, etc.) will see virtually no difference in exposure whether dispersants are used or not. Therefore, organisms that don't ingest tiny oil droplets aren't exposed to different concentrations and so shouldn't be drivers for dispersant decisions (e.g. Herring eggs).


### **General Toxicity Results**

Once dispersant is applied to a spill, three different mixtures might reasonably occur: 1) dispersant, 2) oil, and 3) dispersant/oil mixture; with each exhibiting different toxicities to water column organisms. As discussed above, dispersants are still toxic but have been formulated to be less toxic than the oil they are dispersing. Additionally, operational guidelines target applying dispersants at 1/20th the targeted oil volume so environmentally-relevant dispersant concentrations are a small fraction of the acute oil-chemical risk to water column organisms during dispersant operations. If the threshold of lethality of a particular organism or life stage of an organism is low, then dispersing may create a larger injury zone. In short, the ecological and toxicological risks posed by typical dispersant operations are more closely related to what dispersants do (e.g. put oil into the near surface water column and potentially increase the exposure to certain species) than their own, inherent chemical toxicity.

Another complication in the world of environmental toxicology is the varying sensitivities of different organisms and life stages. Although many California endemic species have been used in toxicity studies involving oil and dispersants (including red abalone, giant kelp, mysids shrimp, Chinook salmon and top smelt) they may not reliably represent all the species of concern found in the Gulf of the Farallones and Cordell Bank (e.g., black abalone, dungeness crab). Without species-specific data, reasonable inferences can often be made from existing data by using appropriate surrogate species.



**Figure 1:** Marine fish Species Sensitivity Distribution (SSD) curve showing LC50s when exposed for 96 hours to chemically dispersed oil using Corexit 9500 LC50 concentrations. LC50 concentrations are given on the x-axis in ug/L of dispersed oil/L of water (equivalent to parts per billion or ppb) and cumulative percentage of test organisms are provided on the y-axis. The CAFE-extrapolated curve is based on data in the 2005 NRC book, Oil Dispersants: Efficacy and Effects.



**Figure 2:** Marine crustacea Species Sensitivity Distribution (SSD) curve showing LC50s when exposed for 96 hours to chemically dispersed oil using Corexit 9500. LC50 concentrations are given on the x-axis in ug/L of dispersed oil/L of water (equivalent to parts per billion or ppb) and cumulative percentage of test organisms are provided on the y-axis. The CAFE-extrapolated curve is based on data in the 2005 NRC book, Oil Dispersants: Efficacy and Effects.

If particular species of concern found in the Gulf of the Farallones or Cordell Bank are not specifically shown in Figures 1 and 2, then a similar or closely related species may serve as an appropriate surrogate. However, there might not be good surrogates for all species or life stages of concern. If the spectrum of species represented reasonably reflects the spectrum of vulnerabilities to dispersed oil, and species of concern might reasonably fall among them, then focusing on dispersed oil concentrations protective of the most sensitive species should be protective of others as well. Collecting this information and having it available in context at the time of a spill is the responsibility of the NOAA Scientific Support Coordinator (SSC) and Applied Response Technology Specialist.

## **Oceanography of North-Central California**

### **Transport along north-central California Coast**

This presentation summarized the oceanic processes that would affect oil transport and fate along California's north-central coast. Introduced concepts that could influence spill trajectory and response decisions included: coastal upwelling, relaxation, 3-D circulation patterns, and the synoptic, seasonal, and inter-annual variables. An important point is that transport of surface oil and subsurface (i.e. dispersed) oil may be different based upon wind and current patterns at the time of the spill. Oil dispersed in deeper water will move with midwater currents, while oil dispersed at the surface will move with the surface currents as influenced by winds and may move onshore. Thus, oil dispersed offshore may very well reduce the potential for oil impacts to nearshore and shallow water species. This may influence which species become the primary dispersant decision drivers. During times of upwelling, it is expected dispersed oil will remain in the upper water column, while during times of downwelling, dispersed oil will be driven deeper into the water column. It is expected some dispersed oil will travel into the nearshore zone during downwelling. Increased coverage of HF radar would facilitate rapid determination of surface currents. Dr. Largier demonstrated that local expertise must be cultivated in order to rapidly inform decision-making processes.

### **Oil fates in San Francisco Bay: Why is it so hard to capture that oil?**

A discussion of the variable factors contributing to difficulties in modeling oil transport and fate in and around SF Bay was included, and the summary message was the system is highly variable both offshore and along SF Bay. Currents within the Bay are dominated by tides, driven by tidal forcing at the Golden Gate, and freshwater inflow. Oil tends to collect in Raccoon Strait and that buoyant material strongly coalesces at convergent zones resulting from buoyancy fronts and shear zones, topographically steered currents and Langmuir circulation. The strong presence of convergence zones in the Bay ensures that oil slicks will turn into linear features, at which point oil control and trajectory prediction is very difficult; though may be advantageous to skimming operations. In addition, neutrally buoyant pollutants and possibly some portion of buoyant pollutants may be subducted below the surface in the shear zones. Dr. Largier indicated the

importance of tackling the problem before the spill transforms into “strings”. He also included a discussion regarding the utility of HFR radar; additional information for which is available at <http://www.ioos.gov/hfradar/welcome.html>.

### ***Fate and Transport of Dispersed Oil***

Mr. Glen Watabayashi provided a summary of NOAA’s modeling capability and expertise in providing oil spill response support, along with first-hand knowledge of the 1984 M/V Puerto Rican incident. The Emergency Response Division, based in Seattle, maintains a suite of modeling tools, which are used in conjunction with GNOME (General NOAA Operational Modeling Environment), a transport model used to predict pollutant trajectories. Effective use of the model in our region could be improved by greater use of access to HF radar data (which is limited by operational funding). Discussion of the 1984 Puerto Rican incident included how unforeseen winds contributed to a shift in the trajectory of the spill towards the Farallones and north to Bodega Bay. Approximately 2,000 gallons of an earlier version of Corexit was in fact used during that incident, though there was disagreement regarding its effectiveness at treating the spilled lube oil. See Breaker and Bratkovich (1993) for more information on the Puerto Rican incident. Mr. Watabayashi closed with a slide showing priorities for improving modeling efforts, while emphasizing that increased data and modeling capacity is not a panacea for ensuring efficient oil spill response. It was pointed out that having water column information that included vertical density structure in the upper 100 meters of water to complement surface HF radar data would help to track and forecast dispersed oil. The speaker also pointed out that unlike Southern California, there is no regional ocean current forecast modeling for Northern California.

### **Biological Resources in the Sanctuaries**

Biological resources were evaluated for the potential negative effects from dispersants ranging from the simplest plankton to birds and mammals and we assembled a list of species of interest drawn from the larger list of species that occur in the region (Appendix IV). Criteria applied to identify species were guided by, but not limited to: 1) significance in the coastal ecosystem as a keystone species or ecosystem driver, 2) federal or state status under endangered species laws, 3) presence of life stages most susceptible to injury or death from dispersed oil, and 4) location of proposed dispersant use in the Gulf of the Farallones. Ultimately, dispersant-use decisions will be guided by the potential percentage impact on the population and recovery time of a species.

Several speakers presented information on invertebrates, fish, birds and mammals of particular concern in the Sanctuary and most likely to be affected by oil or dispersed oil (see appendices for complete notes and presentations). A summary of their findings are presented below:

### **Zooplankton**

Animal larvae within the plankton that are only capable of passively drifting with currents are generally thought to disperse in the water column close to home; replacement of these larvae will depend on their distribution in the area, how broadly this area was impacted, and time of the year. Larvae and plankton capable of independent movement can range over wider areas; impacts to their area may recover more quickly as plankton from non-impacted areas replace them. If larger areas are affected, then it would take longer to re-populate. Given their wide distribution in the Northern California coastal area, most zooplankton populations are generally not likely to be permanently affected by oil spills and are expected to recover. Drake's Bay specifically, and the Gulf of the Farallones region generally, are larval retention zones, and so have the potential to concentrate spilled oil, resulting in longer exposure to oil or dispersed oil. It is likely that a dispersed oil plume generated by an offshore dispersant operation will rapidly be diluted to concentrations not expected to be problematic to species occur in sufficient concentration to affect species within coastal embayments. Larval stages of invertebrates and fish are considered susceptible to oil or dispersants in the water column. However, the effects on certain species may be localized and, given their wide larval distribution, there may not be long-term/regional impacts or population-level effects from local dispersant use.

For many of these species, the adult phase is considered the most important one to protect because they generally experience high mortality rates through the larval and juvenile life stages and therefore protecting the adults producing the next generations of offspring is paramount. However, for some species local or regional effects may have serious consequences for populations. Abalone, for example, need to be in close proximity to each other for successful reproduction. Exposure of adult black abalone to oil may affect the survival and recovery of this endangered species.

### **Invertebrates**

The planktonic stages of Dungeness crab (zoea and megalopa) are present in the GFNMS and CBNMS in Jan-May from mid-water to surface, and the majority of larval stages reside outside of the SF Bay. They are important prey of Chinook salmon and other fish. Dungeness are most vulnerable from Dec-May when larvae are present. Depending on the size of and duration of a chemical spill and subsequent window of dispersant application, a portion of that year class might be lost. It is expected that individual larvae will be lost but population-level effects will be unlikely. Market squid also spawn broadly and have large regional movements and are not considered to be vulnerable to spills or dispersants at the population level.

### **Fish**

Salmon congregate where there are prey species which can be found on temperature and/or salinity fronts. Adult phase of salmon on their landward migration are less susceptible to exposure to dispersed oil due to their swimming speeds and generally rapid movement into San Francisco Bay. Juvenile out-migrating salmon are more

vulnerable to oil and dispersed oil due to increased residency time of the small fish in the GFNMS and CBNMS and their less robust swimming capabilities. Rockfish are found wherever suitable habitat is located in the Sanctuaries and nearby areas. In general for salmon, their main prey species and prey location change seasonally: herring/anchovies=Feb-Mar, in the northern inshore area; euphausiids (primarily *T. spinifera*)=Apr-May, between northern inshore and offshore; rockfish=Mar/Apr-Jun, in the offshore area; and herring/anchovies=Jul-Nov, in the central/southern inshore area. This pattern changes in El Niño years when euphausiids swarm offshore (as occurred in 1986) and salmon follow them). Species such as rockfish that do not move widely are considered more vulnerable locally but are generally found at depths that provide significant dilution for dispersed oil and they would be replaced by movement of animals from adjacent areas. Wide ranging species with large populations such as anchovies were not considered to be vulnerable to spills or dispersants at the population level. There is evidence, though, of the effects of undispersed oil on larval herring that spawn in SF Bay and Tomales Bay.

### **Bird and Mammals**

Birds are the most highly visible and some of the best studied group of species affected directly by oil, both by contact and ingestion. Indirect effects to birds include bioaccumulation of toxins through the food web, exposure to secondary chemicals (dispersants), and destruction of habitat or prey resources. Information we have to date is derived primarily from large spills in California since the early 1970s that have mostly occurred in the winter during the nonbreeding season. Consequently, the most common species that have come onshore, and therefore were documented, include common murre, cormorants, loons/grebes, and shearwaters and fulmars. However, effects to vulnerable offshore species (e.g., ashy storm-petrel, Cassin's auklet) are hard to document. Both species are regionally significant, and storm-petrels are a state species of concern. Marbled murrelets, a federally listed species, occur in the region and nest in Santa Cruz but little is known about where, when or how abundant they are in the region. Some sea birds are attracted to slicks on the water because they look like fish oil slicks. Storm-petrels may be inadvertently attracted to sulfurous crude oil slicks because that particular oil smells like krill (on which they feed) that emit similar compounds. Therefore, reducing surface area of a slick may reduce impact by reducing the "attractiveness" of a slick. There is much information on the potential effects of oiling on waterbirds but little information on the effects of dispersants or dispersed oil on feathers or ingestion at environmentally-realistic concentrations.

For mammals, all breeding species are potentially vulnerable to oil spills because of nursing pups/calves that might ingest oil and because most species congregate during feeding. However, most species range widely and are less likely to be affected depending upon the size, location and season of the spill. Pathways for exposure include inhalation, contact and ingestion. The species most vulnerable to exposure to oil are those that rely on fur for insulation including sea otters and fur seals, species that may be regionally important such as harbor porpoise, and nursing pinnipeds that are

constrained to breeding colonies.

Based on integrated studies such as ACCESS, regional hot spots have been identified where many species of all groups congregate. Taking into account the above information, the complexity of the decision making on the application of dispersants is illustrated by the following scenario. If surface oil circulates in Drakes Bay (as a retention zone), decision makers would have to take into account that 1) any larvae there would have longer exposure to both oil or dispersed oil 2) dispersed oil might reduce potential exposure of water birds to oil floating on the surface and of Dungeness crabs if the oil sinks to the bottom but 3) adult crabs also might be susceptible to dispersed oil depending on how deep dispersed oil sinks and how long it takes for the dispersed oil to decrease in concentration by means of turbulent mixing in the water column

The following discussion on human health issues was not widely discussed with the VSWG but the consensus of the group did feel the topic was worth raising.

### **Human Health Issues**

The broad range of monitoring activities and scientific studies that are still in progress as a result of the exposure of living organisms, including humans, to dispersants and to oil as a consequence of the 2010 Deep Water Horizon oil spill in the Gulf of Mexico are still in progress. Observations by fishermen, scientists, and residents of deformed shrimp, fish with lesions, and increased post-incident mortality of baby dolphins continue to be compiled in that region. There are functional differences between evaluations of acute and chronic exposure, and acute and chronic effects, with respect to both oil and to dispersant components, and some important lessons remain as follow-up studies of the Gulf of Mexico spill are completed.

One of any Marine Sanctuary's most valued resources is its people, including Site staff, affiliated partner agencies, and the general public served by the Site. Protection of human health and public safety is a paramount priority for the Sanctuary Program at all times. OSHA guidelines require training of sanctuary staff who are required to wear appropriate protective clothing in any response-related situations where they may come into contact with petroleum products or dispersants, and to utilize individual respiratory protection where indicated. Children, women who are pregnant, and those with compromised immune systems, should not be allowed near spill response or oiled wildlife recovery situations. The public must be fully advised to avoid any oil spill unless they are HAZMAT trained and deployed under the appropriate level of supervision.

## Recommendations

---

### General Science and Research Recommendations

1. Continue to follow and support research on the effects and impacts of alkylated PAHs in dispersed oil on fish and invertebrate egg and larvae.
2. Actively support the research and development of “next generation” biodegradable dispersants and alternative products for oil spill response. Consider establishing a policy that requires the use of alternatives to existing dispersants within the GFNMS and CBNMS.
3. Complete a review of the existing literature and identify data gaps on the status of marine life within the Greater Gulf of the Farallones Ecosystem, particularly during the winter. Identify opportunities for research on species of concern that would be affected by oil spills and dispersants in the GFNMS and CBNMS during winter.
4. Support NOAA and State research on the reproductive success and behavioral effects (spawning, foraging, predator avoidance) for black/red abalone exposed to dispersed oil. Specific research on the potential for behavioral responses from exposure to dispersed oil (narcosis) is needed.
5. Support research on the effects of dispersed oil on critical or surrogate species that represent important commercial and public trust resources in the GFNMS and CBNMS. Targeted research should include adult and juvenile Dungeness crab, and ESA listed Black Abalone and the potential impacts (short and long-term) to the habitats that support these resources.
6. Identify existing sources of real-time data feeds on surface and subsurface currents on the GFNMS and CBNMS (e.g. NOAA Data Buoys, HR Radar). Support the placement of an HR radar antenna on Southeast Farallon Island to close the existing radar shadow.
7. Identify current or published research on the effects of inhalation and dermal exposure of dispersed and non-dispersed oil on birds and marine mammals.
8. Identify the seasons and species that use GFNMS and CBNMS in substantive numbers where an oil spill and/or dispersed oil could have significant long-term impacts on the viability of the population (e.g. Ashy Storm-petrels).
9. Support research that includes:
  - a) Standardization of dispersant toxicity studies for inter-comparability,
  - b) Maximizing dispersant efficacy while minimizing potential toxicity, and

- c) Filling data gaps on:
  - 1) Feather and fur wetting effects by environmentally realistic concentrations of dispersed oil, and
  - 2) Toxicity testing of species of concern and how they relate to surrogate species and species sensitivity curves.
- 10. Support research to find more effective seagoing and coastal oil spill containment and sorbent booms, skimmers, separators, and “oil mop” types of petroleum recovery devices for use in GFNMS and CBNMS.

### **General Education and Outreach Recommendations**

1. The GFNMS and CBNMS Superintendents need to establish an annual coordination meeting with NOAA ERD and ARD, EPA, USCG and OSPR on coordinating the San Francisco Bay-Delta Contingency Plan pre-spill planning with the Sanctuary roles/response coordination.
2. The GFNMS and CBNMS Superintendents in coordination with OSPR need to develop an oil spill and response outreach plan for county and local governments that border the GFNMS in order to foster communication and awareness and to establish pre-spill working relationships.
3. The GFNMS and CBNMS Superintendents need to schedule an annual USCG and NOAA OR&R briefing at the joint SAC Meeting to provide SAC members updates on spill modeling, cleanup technologies, dispersants exposure research, non-toxic dispersant development or any emerging news on oil spill containment and response such as gelling agents, emulsion breakers, improved chemical spreading additives to enhance physical mixing/dispersant effectiveness.

### **General Policy and Management Recommendations**

1. Seek funding to complete the SW ERMA placing a priority on the GFNMS and CBNMS and in the process of building data sets. Identify the highest priority/most sensitive species at risk during an oil spill for inclusion in the SW ERMA.
2. Working with the USCG, EPA and OSPR, develop a standing policy that provides for using commercial fishermen in response and clean-up which takes advantage of local knowledge and expertise to most effectively deploy response assets.
3. The GFNMS and CBNMS Superintendents need to support the development of a specialized NEBA within the Sanctuaries that focuses on specific resources and/or physical events such as seasonal upwelling, and sensitive habitats that support nearshore and subtidal species that are known to be highly sensitive to oil and/or dispersed oil (e.g. Dungeness crab, black and red abalone).

## Specific Sanctuary Recommendations

1. Given the Superintendent's role is advisory/consultative to the RRT, the SAC recommends a precautionary approach to any incident response technology. America's National Marine Sanctuaries are "Special Ocean Places", worthy of special national recognition and protection. Any oil spill response decisions in the Sanctuaries waters will require a higher burden of proof of compelling need given the high resource productivity and sensitivity.
2. The Superintendents need to consider a policy of no-aerial spraying area within one mile of the Farallon Islands. If warranted boats would be authorized within one mile to apply dispersant in water >60' deep. In terms of other mainland coastal rookeries, haul-out sites and areas identified as sensitive habitats should follow the provisions of the Wildlife Response Plan.
3. To assist the Superintendents in making decisions on the application of dispersants attention needs to be given to the Sensitive Species Matrix (Appendix V), and that the Matrix is modified as new science-based information is obtained.
4. Provisions need to be made to review additional data collection needs and updating of the Sensitive Species Matrix (Appendix V) should the boundaries of the GFNMS and CBNMS change.
5. It is suggested that the Sanctuary Superintendents request that the appropriate public health entities (e.g. NIH, EPA, Public Health Departments, etc.) provide information regarding the human health effects of oil, dispersants, and dispersed oil on responders and general public. Sanctuary Superintendents consider this information in the deployment of Sanctuary staff and resources while actively supporting the research and development of alternative products for oil spill response (General Science and Research Recommendation #2).

## Appendix I: Meeting Agendas

---

All meetings held at PRBO Conservation Science, 3820 Cypress Drive Petaluma, CA

### June 20, 2011

Review WG process, ground rules, and timeline and presentation listed below:

1. **Irina Kogan**, Spill Response Structure and Decision Making Authority
2. **Yvonne Addassi**, Response Technology Overview and Dispersants 101

### August 30, 2011

Review oil spill response decision making process as listed below:

1. **Yvonne Addassi**, NEBA overview
2. **Jordan Stout**, Spill Scenario and Trajectory
3. **Ellen Faurot-Daniels**, Simulation of Process for Dispersant Use/Non-use Recommendation Development to the Unified Command

### February 2, 2012

Review toxicology of oil and dispersed oil with the speakers listed below:

1. **Ron Tjeerdema**, Toxicological effects of dispersed (and not dispersed) oil on 1) Salmon smolts and adults, 2) Fish larvae (top smelt), 3) Abalone larvae and adults, 4) Zooplankton species (mysids), 5) Kelp, and persistence of dispersants in the environment and bioaccumulation COREXIT ingredients and DOSS (dioctyl sodium sulfosuccinate).
2. **Gary Cherr**, Bodega Marine Lab, UC Davis Oil and Embryos Do Not Mix: Impacts of the Cosco Busan Bunker Fuel Oil Spill on Pacific Herring
3. **Alan Mearns**, Marine Environmental Tradeoffs of Dispersant Operations: Knowns, Unknowns and Dealing with Uncertainty, Toxicological effects of dispersed (and not dispersed) oil on 1) Rockfish, 2) Crab, 3) Oyster, mussels and other mollusks (not abalone), 4) Copepods and krill in water column, 5) Top predators (seabirds, pinnipeds, whales, etc).

### March 8, 2012

Review bio-resources in the sanctuaries with the speakers listed below:

1. **Pete Kalvass**, Crab biology and population dynamics, spatial and temporal patterns in distribution, phenology, important times in their development most critical for spill response.
2. **Pete Adams**, Salmon Foraging areas, biology and population dynamics, spatial and temporal patterns in distribution, phenology, important times in their development most critical for spill response.

3. **Pete Warzybok** Birds and mammals, biology and population dynamics, spatial and temporal patterns in distribution, phenology, important times in their development most critical for spill response.

#### **April 12, 2012**

Review oceanography in the sanctuaries with the speakers listed below:

1. **John Largier**, Physical oceanography and how local conditions may affect the fate of non-dispersed and dispersed oil in the greater Gulf of the Farallones (currents?). Also currents and larval transport.
2. **Toby Garfield**, Physical oceanography and how local conditions may affect the fate of non-dispersed and dispersed oil in San Francisco Bay due to tides and currents.
3. **Al Venosa**, How deeply dispersed oil will sink? How quickly dispersed oil/dispersant breaks down? How quickly water column concentrations of oil with dispersant decrease to zero? What are the protocols for monitoring the effects of chemically dispersed oil on water column biota?
4. **Glen Watabayashi**, How you predict movement of dispersed oil? What are the parameters you use to predict how quickly the dispersed oil will break down; what can we expect in this region? How confident is NOAA about their predictions for this area? 4) how deeply would dispersed oil sink in this area, what are the local controls?

#### **May 3, 2012**

Review, science, outreach, and policy recommendations compiled to date.

#### **May 10, 2012**

Review toxicology section and other major changes needed to background document. Review recommendations to date and draft list of sensitive species.

## Appendix II: Working Group Member Biographies

---

**Yvonne Addassi** is a bright scientist from OSPR who didn't prove a bio.

**Sarah G. Allen** is program lead for the Coast and Oceans Program of the National Park Service, Pacific West Region. She received her B.S., M.S. and Ph.D. from the University of California, Berkeley. She has been studying marine birds and mammals for more than 35 years, mostly in California. She has served on the Scientific Peer Review Panel of the San Francisco International Airport, the Governing Council of the Central and Northern California Ocean Observing System (CeNCOOS- <http://www.cencoos.org/>), NOAA Subtidal Habitat working group for San Francisco Bay, and the Science Advisory Team of north-central California Marine Life Protection Act Master Plan.

**Richard Charter** has worked for the past 34 years on ocean protection issues, including marine spatial planning, congressional liaison activities in support of conservation outcomes, and preventing and mitigating industrial impacts on marine habitats. Richard works with local and state elected officials, the fishing community, and regional and national NGO interests to secure advances in sustainable management of ocean ecosystems, marine and estuarine resource restoration projects, and protection of ocean-based regional economies and public health in the context of offshore oil and gas drilling issues. As Co-Chair of the National Outer Continental Shelf (OCS) Coalition, Richard was involved in initiating and maintaining the twenty-seven-year congressional moratorium on offshore oil and gas leasing which has thus far prevented new drilling along the U.S. West Coast, the Atlantic Coast, and Florida's Gulf Coast, as well as in Alaska's Bristol Bay. Richard also coordinated the local government support that helped to bring about the creation of the Gulf of the Farallones, Cordell Bank, Channel Islands, and Monterey Bay National Marine Sanctuaries. Richard serves on the Department of Energy's Methane Hydrates Advisory Committee FACA and is serving in his second term as chair of the Gulf of the Farallones Sanctuary Advisory Council.

**Ellen Faurot-Daniels** has a BA in Biological Sciences and an MS in Marine Science. Past work includes various CDFG projects (1979-90), environmental group Science Director (1992-97), university research assistant (1997-98), Oiled Wildlife Care Network (1997-98), and supervisor of the California Coastal Commission Oil Spill Program (1998-2009). She has been actively involved in oil spill prevention and response planning since 1992, was a working group member for various NMS planning teams, and was a stakeholder on two MLPA workgroups. One of her several oil spill responses included the Deepwater Horizon. She has been with CDFG-OSPR since 2009, coordinating the licensing and use of oil spill cleanup agents, the development of statewide policies for the use of applied response technologies, and serving as technical specialist for both ART and fishery closures during oil spills.

**Joe Dillon** is the Southwest Regional Water Quality Coordinator for NOAA's National

Marine Fisheries Service (NMFS). He has been with NMFS since 1999 serving in a number of capacities involving water quality, toxicology and fate and transport of pollutants. He was designated the NMFS oil spill responder for California in 2002 and continues to be the main NMFS point of contact for these issues from Monterey County north to the Oregon border (inclusive).

**Barbara Emley** started commercial fishing with her husband (Larry Collins) in 1985. She has fished for Salmon, Crab, Albacore and Rockfish using hook and line and traps. Barbara represents commercial fishing on the GFNMS SAC. She also is on the board of the Institute for Fisheries Resources (IFR); Pacific Coast Federation of Fishermen's Associations (PCFFA); and the California Salmon Council. Barbara is the public Commissioner for California on the Pacific States Marine Fisheries Commission (PSMFC). She and her husband worked together to form the San Francisco Community Fishing Association which just finished its first year of operation.

**Jaime Jahncke** joined PRBO in 2004. He received his Ph.D. in Biological Sciences from the University of California Irvine (2004). Jaime's doctorate research focused on how physical processes associated with coastal waters affect the abundance and distribution of marine birds in Peru and Alaska. Jaime's current research focuses on the spatial and temporal relationships between oceanographic processes, zooplankton, and marine birds and mammals in the region surrounding Cordell Bank and the Gulf of the Farallones. Jaime is the lead Principal Investigator for the Applied California Current Ecosystem Studies (ACCESS) Partnership between PRBO, NOAA's Cordell Bank and Gulf of the Farallones National Marine Sanctuaries and several agencies and academic institutions. He currently participates on both Cordell Bank and Gulf of the Farallones National Marine Sanctuaries Sanctuary Advisory Councils (Primary for Cordell Bank and alternate for Gulf of the Farallones).

**Gerry McChesney** has a B.A. in Biology with a focus in Marine Sciences, U.C. Santa Cruz, 1988. M.S. in Biology (Conservation Biology), Sacramento State University, 1997. He has been studying seabirds in California since 1986, focusing on breeding population surveys and breeding biology but also has conducted studies of at-sea distribution. Wildlife Biologist for Humboldt State University from 1989-2002. Has worked for the U.S. Fish and Wildlife Service, San Francisco Bay National Wildlife Refuge Complex, since 2002. Since 2002, has managed the Refuge's seabird restoration program and since 2008 has also managed the Farallon National Wildlife Refuge.

**Patrick Rutten** has worked for NOAA for 35 years in ocean, coastal/estuarine research, and management positions within NOAA and NMFS. His career has focused on California fisheries and coastal habitat management. After 21 years of sea service in the Gulf of Alaska, Bering Sea, eastern Pacific and Hawaii with the NOAA Corps he retired in 1995 at the rank of Commander to assume the Central California Field Supervisor for NMFS, Protected Resources Division. In 2004 he took a new position with the NOAA Restoration Center as Southwest Field Supervisor for California and the Pacific Islands

administering the Community-based Restoration Program and Damage Assessment and Remediation Program. Mr. Rutten has a B.S., Marine Biology, Cal Poly San Luis Obispo, CA, and a M.S. Management, Naval Postgraduate School, Monterey, CA.

**Deb Self** is Executive Director of San Francisco Baykeeper, an organization dedicated to protecting San Francisco Bay from pollution. Deb [d1] is Vice Chair of California's Office of Spill Preparedness and Response Technical Advisory Committee and is a member of the San Francisco Harbor Safety Committee. Deb also serves as Chair of the Waterkeeper Alliance, an international movement of on-the-water advocates for fishable, swimmable, drinkable waters, and is a Board member of the California Coastkeeper Alliance, the statewide affiliation of Waterkeeper organizations. Deb has a Masters in sociology focused on environmental justice and holds a bachelors degree in geology.

**Jordan Stout** currently serves as the NOAA Scientific Support Coordinator (SSC) here in California where he provides scientific & technical support to USCG & EPA for oil spills & hazmat releases. He has been involved in many significant incidents/responses in California and throughout the nation, including: *SS Montebello* assessment, Japanese Tsunami response, *MODU Deepwater Horizon (MC252)*, *T/V Dubai Star*, *M/V Selendang Ayu*, *M/V Cosco Busan*, Sacramento River Humpback Whales, Hurricane *Katrina*, and numerous others. Jordan also serves as NOAA's representative on RRT-9 and the MEXUS-PAC Joint Response Team. Jordan has prior work experience with the USFWS' Environmental Contaminants Program in Alaska and with Miami-Dade County's Department of Environmental Resources Management. He holds a Master's of Environmental Management from Duke University's Nicholas School of the Environment (Focus: Environmental Toxicology & Risk Assessment) and a BS from the University of Miami (Double-major: Marine Science & Biology; Minor: Chemistry).

**Bob Wilson** is active in a number of environmental organizations. He is the former Chair of the Farallones Marine Sanctuary Association board, an acting executive director and is currently CFO. He has been an active Beachwatch and SEALS volunteer. He is on the GFNMS Sanctuary Advisory Council. He is a director emeritus of The Marine Mammal Center, is currently Policy Liaison for TMMC and is an animal care and rescue volunteer. Bob is on the audit committee of the Desert Tortoise Preserve Committee. He is also on the advisory board of the Snow Leopard Conservancy. He is an attorney and retired from the Federal government.

**Lieutenant Commander Blanca Rosas** reported to her current position as the Chief of Incident Management Division (IMD) at Sector San Francisco in July of 2011, having completed a staff assignment as Chief of the Officers Promotions Section at Personnel Service Center (OPM-1) in Arlington, VA. LCDR Rosas joined the Coast Guard through the College Student Pre-Commissioning Initiative program in 1997. She acquired her commission as an Ensign from Officer Candidate School in February of 2000. Her prior positions include Marine Environmental Protection Chief

at Sector Delaware Bay, Philadelphia and Contingency Planner / IMD Deputy Chief at Sector San Juan, Puerto Rico. LCDR Rosas has a B. S. in Computer Science from the University of Puerto Rico. Her military awards include the Coast Guard Commendation medal, two Achievement medals and various team awards.

**Michael Carver** is the Deputy Superintendent for Cordell Bank National Marine Sanctuary. Michael has been with Cordell Bank National Marine Sanctuary since 2000. Michael started the sanctuary's monthly at sea monitoring program and managed it for several years before moving on. Michael's responsibilities include overseeing enforcement, permitting, planning, and management actions to address threats to the Sanctuary. Michael also provides engineering support for sanctuary field operations, and serves as the staff lead on emergency response issues. In addition, Michael coordinates annual budget planning and execution, interagency agreements, manages contracts, and works closely with the sanctuary superintendent to ensure smooth operation of the sanctuary.

**Irina Kogan** is the Oil Spill Response Coordinator and Permit Coordinator at the Gulf of the Farallones National Marine Sanctuary. She has experience responding to oil spills and an academic background in Geology and Geochemistry. Irina's spill response and preparedness activities include representing the GFNMS at ACP and RRT meetings and ensuring GFNMS staff are prepared to participate in spill response activities affecting the sanctuaries.

---

## Appendix III: Invited Speaker Biographies

---

**Dr. Ron Tjeerdema** received his PhD in Pharmacology & Toxicology from UC Davis. Initially a faculty member in the Department of Chemistry & Biochemistry at UC Santa Cruz, he is now Professor and Chair of the Department of Environmental Toxicology at UCD. He also currently holds the Donald G. Crosby Endowed Chair in Environmental Chemistry, and is certified in General Toxicology by the American Board of Toxicology. In over 25 years, Dr. Tjeerdema has attracted some \$30 million in extramural research support and published in excess of 200 peer-reviewed research articles. His areas of expertise range from chemical fate in the environment, sensitive lifestage bioassays and biochemical mechanisms of toxicity. He has worked extensively with pesticides, marine planktonic toxins, and petroleum hydrocarbons and dispersants. Due to his extensive work with dispersants and dispersed oil, Dr. Tjeerdema served on the NOAA panel that recommended dispersant injection during the Gulf Oil Spill. He has since served on panels regarding the Gulf Spill for NOAA, EPA and NCEAS, and provided testimony to both the House of Representatives Subcommittee on Natural Resources and the President's Oil Spill Commission; he continues as an advisor to the California Office of Spill Prevention & Response. Dr. Tjeerdema is an Editor-in-Chief of Aquatic Toxicology, and serves on the editorial boards of the Reviews of Environmental Contamination & Toxicology, Marine Pollution Bulletin and Bulletin of Environmental Contamination & Toxicology.

**Dr. Gary Cherr** received his Ph.D. from the University of California Davis, was an NIH postdoctoral fellow, and has worked in reproductive and developmental toxicology for over 25 years. Dr. Cherr is Professor of Environmental Toxicology and Nutrition at the University of California Davis, and is currently the Director of the University of California Davis' Bodega Marine Laboratory. His laboratory studies stressor impacts on marine invertebrates and vertebrates, and embryo defenses to chemical stressors. Dr. Cherr's laboratory has investigated the impacts of salinity stress and petroleum hydrocarbons on fish embryos in S.F. Bay for over 20 years. His group was the first to show that creosote-treated wood pier pilings were toxic to herring embryos. Dr. Cherr serves on the Exxon Valdez Oil Spill Trustee Council's Science Panel, is a member of the National Center for Ecological Analysis and Synthesis' Gulf Oil Spill Working Group, and his laboratory participated in the NRDA for the 2007 Cosco Busan fuel oil spill in San Francisco Bay.

**Dr. Alan Mearns** is a marine ecologist and Senior Staff Scientist with NOAA's national Emergency Response and Division (ERD) in Seattle, Washington. He supports NOAA's regional Scientific Support Coordinators (SSC's) and U.S. Coast Guard Sector Offices during spills of oil and hazardous materials. Alan received his B.Sc. and M.A. degrees in Biology from California State University at Long Beach, and his Ph.D. in Fisheries from the University of Washington. During the 1970's, he was Leader of the Biology Division at the Southern California Coastal Water Research Project (SCCWRP) where his team

pioneered studies on the effects and ecological tradeoffs of wastewater treatment for large ocean sewage outfalls. He joined NOAA in Seattle in 1980 serving as Ecologist for the Puget Sound MESA (Marine EcoSystem Assessment) Program. During the mid- and late 1980's he helped develop the NOAA National Status and Trends Program and continues to support the National Mussel Watch. In 1989 Alan participated in the first assessment surveys of the Exxon Valdez Oil Spill in Prince William Sound, then joined the ERD team in Seattle. Alan has provided support to NOAA and the US Coast Guard for dozens of major spill responses around the US and internationally. He provided 24/7 support on the 2010 Deepwater Horizon spill including focus on assessment of dispersant operation effectiveness and effects. Alan was Vice-Chair of the Technical Advisory Committee for the Santa Monica Bay Restoration Project, and a member of the Committee of Science Advisors for the San Francisco Estuary Institute (SFEI). In 1999 and again in 2004 Alan had the privilege of serving on both the 5- and 10-year science advisory board review committees for the San Francisco Bay Toxic Substances Monitoring Program conducted by SFEI. He also provided technical support on wastewater management for the Alaska Cruise Ship Initiative. Alan is a member of the Science and Technical Committee of the Oil Spill Recovery Institute (OSRI) in Cordova, Alaska. Dr. Mearns received the 1985 Biology Alumnus of the Year Award from California State University at Long Beach, a 1992 Silver Medal from the US Department of Commerce for work on the Exxon Valdez oil spill response and is listed in American Men and Women of Science.

**Pete Kalvass** - California Department of Fish and Game, Marine Region

**Pete Adams**, is the recently retired Fisheries Investigation Chief of the Southwest Fisheries Science Center, National Marine Fisheries Service, Santa Cruz, California, U.S.A., where he had overall responsibility for providing scientific advice on Southwest salmon and groundfish, the Endangered Species Act (ESA), and harvest management. His personal research focus has been on: (1) assessing population viability under the ESA, (2) harvest advice including the use of life history theory, (3) sample survey design, and (4) methods to communicate the level of uncertainty associated with estimates to decision makers.

**Pete Warzybok**, joined PRBO a volunteer seabird research assistant on the Farallon Islands in the spring of 2000, and was hired as a seabird biologist the following year. Prior to coming to PRBO, he worked on a waterfowl management project in suburban NY, and on seabird monitoring projects with USGS in Alaska and USFWS in Maine. Pete received his B.S. in biology from the State University of New York at Purchase in 1996, where his undergraduate research focused on geographic variation in the song of Brown-headed Cowbirds. His current research interests include diet, prey availability, and ecosystem variability and their effects on the breeding success and population dynamics of Farallon seabirds.

**Dr. John Largier** John Largier is Professor of Coastal Oceanography at the University of

California Davis (UCD), resident at Bodega Marine Laboratory. Prior to 2004, he was Research Oceanographer at Scripps Institution of Oceanography. He has also held positions at the University of Cape Town and the National Research Institute for Oceanology (CSIR) in South Africa. His research, teaching and public service is motivated by contemporary environmental issues and centered on the role of transport in ocean, bay, nearshore and estuarine waters. His work has addressed transport of plankton, larvae, contaminants, pathogens, heat, salt, nutrients, dissolved oxygen, and sediment – and he places this work in the context of issues as diverse as marine reserves, fisheries, mariculture, beach pollution, wastewater discharge, wildlife health, desalination, river plumes, coastal power plants, kelp forests, wetlands, marine mining, coastal zone management and impacts of coastal development. At UCD he heads the 16-person Coastal Oceanography Group. Dr Largier is a leader in developing the field of “environmental oceanography” through linking traditional oceanographic study to critical environmental issues. Dr Largier serves on the Science Advisory Team for the California Marine Life Protection Act (MLPA), the Governing Council for CeNCOOS (Central and Northern California Ocean Observing System), the Sanctuary Advisory Committee for the Gulf of Farallones, and several other advisory boards. He is president of the California Estuarine Research Society. In 2002-2004, Dr Largier played a significant role in advising the state on beach pollution and in the late 1990’s, he played a key role in developing the knowledge foundation for the new coastal zone management policy in South Africa. He is an Aldo Leopold Leadership Fellow. Following undergraduate studies in Maths and Physics, he obtained a Ph.D. in Oceanography from the University of Cape Town (South Africa) in 1987.

**Toby Garfield**, my research focus is ocean current circulation along the continental margin, the region from the shore out to and beyond the continental shelves and slopes. Using both traditional tools and the new technologies of satellites and autonomous sampling vehicles I have studied the ocean circulation in the Gulf of Maine, the Brazil Current at tropical latitudes and the shelf and slope circulation between Pt. Sur and Bodega California. In addition I have conducted two studies in the Gulf of the Farallones and studied the California Undercurrent, a current that flows poleward, carrying subtropical water north into the North Pacific. Presently I am working with the CA Coastal Conservancy to establish a network of surface current monitoring instruments that will measure the coastal ocean circulation along the whole California coast and provide maps and data on the web in near real time

**Albert D. Venosa**, Director of Land Remediation and Pollution Control Division in EPA’s National Risk Management Research Laboratory, Cincinnati, Ohio. For the past 20 years, Al has led EPA’s oil spill research and development program to conduct basic and applied research in both the laboratory and the field in the area of spill response technology development. Al was an EPA team leader in the Exxon Valdez bioremediation project in 1989 and 1990. Al also conceived and led an important controlled oil spill project on the shoreline of Delaware Bay in 1994, which demonstrated statistically that bioremediation with simple inorganic nutrients enhances

the biodegradation rate of crude oil on a marine shoreline compared to natural attenuation without amendments. Al repeated a similar experiment in 1999 on a Quebec freshwater wetland and again in 2001 on a Nova Scotia salt marsh in collaboration with our Canadian government partners. In addition to those field studies, Al led a research team in developing laboratory protocols to test the effectiveness of commercial bioremediation agents and chemical dispersant products for use in treating oil spills. Al I have conceived and led numerous other studies to understand how best to respond to and mitigate oil spills on land.

**Glen Watabayashi**, is the lead physical oceanographer for NOAA's Emergency Response Division (ERD) of the Office of Response and Restoration (OR&R). Glen started working for NOAA on modeling currents in 1974 while at the U. of W. School of Oceanography. Glen has worked on a number of high profile oil spills including: IXTOC ('79), Exxon Valdez ('89), First Gulf War('91), Deep Water Horizon ('10). Glen has also lent his expertise to other projects including the search for black box from Korean Airlines 007('83), the debris field from JFK Jr. airplane crash ('99) and the debris field from Japan tsunami debris ('12). In addition to this work Glen has also made significant computer code contributions for: ALOHA (1,2,3,4,5) OSSM (original oil spill model) ADIOS I. Glen, grew up in Honolulu, went to Columbia University for a BA and received his masters in physical oceanography from University of Washington.

## Appendix IV: List of Marine Species of GFNMS and CBNMS

---

### Important definitions

**Sensitivity to oil:** Species likely to have strong negative population effects due to a localized oiling event

**More benefit from dispersants:** Species likely to benefit from removing spilled oil from the surface of the ocean, or from reducing the amount of oil likely to hit the shoreline

**More harm from dispersants:** Species likely to be affected by the oil dispersed in the water column

**Endangered Species Listing:** Species currently included in the Federal endangered species list and in the California lists of species of special concern

**Breeds locally:** Species that breed within the Sanctuaries

**Reason:** The specific reason why the working group decided a species was sensitive to oil and would (or would not) benefit from dispersants

### VERTEBRATES

#### Mammals

Scientific Name	Common Name	Sensitivity to oil	More Benefit from Dispersant	More Harm from Dispersant	ESA listing	Breeds locally	Reason
<i>Balaenoptera musculus</i>	Blue Whale	Low	yes	no	E	no	Potentially affected by oil fumes, direct and indirect effects of oil ingestion, oiling of baleen
<i>Balaenoptera physalus</i>	Fin Whale	Low	yes	no	E	no	Potentially affected by oil fumes, direct and indirect effects of oil ingestion, oiling of baleen
<i>Balaenoptera borealis</i>	Sei Whale	Low	yes	no	E	no	Potentially affected by oil fumes, direct and indirect effects of oil ingestion, oiling of baleen
<i>Balaenoptera acutorostrata</i>	Minke Whale	Low	yes	no		yes	Potentially affected by oil fumes, direct and indirect effects of oil ingestion, oiling of baleen
<i>Megaptera novaeangliae</i>	Humpback Whale	Low	yes	no	E	no	Potentially affected by oil fumes, direct and indirect effects of oil ingestion, oiling of baleen
<i>Eschrichtius robustus</i>	Gray Whale	Low	yes	no	D	yes	Potentially affected by oil fumes, direct and indirect effects of oil ingestion,

<i>Eubalaena glacialis</i>	Northern Right Whale	Low	yes	no	E	no	oiling of baleen Potentially affected by oil fumes, direct and indirect effects of oil ingestion, oiling of baleen
<i>Phocoena phocoena</i>	Harbor Porpoise	Medium	yes	no		yes	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Phocoenoides dalli</i>	Dall's Porpoise	Low	yes	no		yes	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Lagenorhynchus obliquidens</i>	Pacific White-sided Dolphin	Low	yes	no		yes	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Lissodelphis borealis</i>	Northern Right Whale Dolphin	Low	yes	no		yes	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Delphinus delphis</i>	Short-beaked Common Dolphin	Low	yes	no		no	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Delphinus capensis</i>	Long-beaked Common Dolphin	Low	yes	no		no	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Tursiops truncatus</i>	Bottlenose Dolphin	Low	yes	no		yes	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Stenella coeruleoalba</i>	Striped Dolphin	Low	yes	no		no	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Stenella attenuata</i>	Spotted Dolphin	Low	yes	no		no	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Steno bredanensis</i>	Rough-toothed Dolphin	Low	yes	no		no	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Grampus griseus</i>	Risso's Dolphin	Low	yes	no		no	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Orcinus orca</i>	Killer Whale	Low	yes	no	E	no	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Globicephala macrorhynchus</i>	Short-finned Pilot Whale	Low	yes	no		no	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Physeter macrocephalus</i>	Sperm Whale	Low	yes	no	E	no	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Kogia breviceps</i>	Pygmy Sperm Whale	Low	yes	no		no	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Kogia simus</i>	Dwarf Sperm Whale	Low	yes	no		no	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Ziphius cavirostris</i>	Cuvier's Beaked Whale	Low	yes	no		no	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Berardius bairdii</i>	Baird's Beaked Whale	Low	yes	no		no	Potentially affected by oil fumes, direct

<i>Mesoplodon calrhubbsi</i>	Hubb's Beaked Whale	Low	yes	no		no	and indirect effects of oil ingestion Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Mesoplodon densirostris</i>	Blainsville's Beaked Whale	Low	yes	no		no	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Mesoplodon stejnegeri</i>	Steineger's Beaked Whale	Low	yes	no		no	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Eumetopius jubatus</i>	Steller Sea Lion	Medium	yes	no	T	yes	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Zalophus californianus</i>	California Sea Lion	Medium	yes	no		yes	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Callorhinus ursinus</i>	Northern Fur Seal	High	yes	no		yes	Oil effects on insulation, direct and indirect effects of oil ingestion
<i>Arctocephalus townsendi</i>	Guadalupe Fur Seal	High	yes	no	T	no	Oil effects on insulation, direct and indirect effects of oil ingestion
<i>Mirounga angustirostris</i>	Northern Elephant Seal	Medium	yes	no		yes	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Phoca vitulina</i>	Harbor Seal	Medium	yes	no		yes	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Enhydra lutris</i>	Sea Otter	High	yes	no		yes	Oil effects on insulation, direct and indirect effects of oil ingestion
<i>Lantra canadensis</i>	River Otter	Low	yes	no		yes	Estuarine species and studies shown low sensitivity

---

### Birds

Scientific Name	Common Name	Sensitivity to oil	More Benefit from Dispersant	More Harm from Dispersant	ESA listing	Breeds locally	Reason
<i>Gavia stellata</i>	Red-throated Loon	High	yes	no		no	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Gavia pacifica</i>	Pacific Loon	High	yes	no		no	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Gavia immer</i>	Common Loon	High	yes	no	SC	no	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Gavia adamsii</i>	Yellow-billed Loon	High	yes	no		no	Typical feather and ingestion effects,

<i>Podilymbus podiceps</i>	Pied-billed Grebe	High	yes	no		no	experience shows frequent oiling during oil spills
<i>Podiceps auritus</i>	Horned Grebe	High	yes	no		no	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Podiceps grisegena</i>	Red-necked Grebe	High	yes	no		no	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Podiceps nigricollis</i>	Eared Grebe	High	yes	no		no	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Aechmophorus occidentalis</i>	Western Grebe	High	yes	no		no	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Aechmophorus clarkii</i>	Clark's Grebe	High	yes	no		no	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Phoebastria immutabilis</i>	Laysan Albatross	High	yes	no		yes	Oil effects on insulation, inhalation, direct and indirect effects of oil ingestion
<i>Phoebastria nigripes</i>	Black-footed Albatross	High	yes	no		yes	Oil effects on insulation, inhalation, direct and indirect effects of oil ingestion
<i>Phoebastria albatrus</i>	Short-tailed Albatross	High	yes	no	E	no	Oil effects on insulation, inhalation, direct and indirect effects of oil ingestion
<i>Fulmarus glacialis</i>	Northern Fulmar	High	yes	no		no	Oil effects on insulation, inhalation, direct and indirect effects of oil ingestion
<i>Pterodroma ultima</i>	Murphy's Petrel	High	yes	no		no	Oil effects on insulation, inhalation, direct and indirect effects of oil ingestion
<i>Pterodroma inexpectata</i>	Mottled Petrel	High	yes	no		no	Oil effects on insulation, inhalation, direct and indirect effects of oil ingestion
<i>Pterodroma phaeopygia</i>	Dark-rumped Petrel	High	yes	no	E	no	Oil effects on insulation, inhalation,

<i>Puffinus creatopus</i>	Pink-footed Shearwater	High	yes	no	no	no	direct and indirect effects of oil ingestion Oil effects on insulation, inhalation, direct and indirect effects of oil ingestion
<i>Puffinus carneipes</i>	Flesh-footed Shearwater	High	yes	no	no	no	Oil effects on insulation, inhalation, direct and indirect effects of oil ingestion
<i>Puffinus bulleri</i>	Buller's Shearwater	High	yes	no	no	no	Oil effects on insulation, inhalation, direct and indirect effects of oil ingestion
<i>Puffinus griseus</i>	Sooty Shearwater	High	yes	no	no	no	Oil effects on insulation, inhalation, direct and indirect effects of oil ingestion
<i>Puffinus tenuirostris</i>	Short-tailed Shearwater	High	yes	no	no	no	Oil effects on insulation, inhalation, direct and indirect effects of oil ingestion
<i>Puffinus puffinus</i>	Manx Shearwater	High	yes	no	no	no	Oil effects on insulation, inhalation, direct and indirect effects of oil ingestion
<i>Puffinus opisthomelas</i>	Black-vented Shearwater	High	yes	no	no	no	Oil effects on insulation, inhalation, direct and indirect effects of oil ingestion
<i>Oceanites oceanicus</i>	Wilson's Storm-Petrel	High	yes	no	no	no	Oil effects on insulation, inhalation, direct and indirect effects of oil ingestion
<i>Oceanodroma furcata</i>	Fork-tailed Storm-Petrel	High	yes	no	no	no	Oil effects on insulation, inhalation, direct and indirect effects of oil ingestion
<i>Oceanodroma leucorhoa</i>	Leach's Storm-Petrel	High	yes	no	yes	yes	Oil effects on insulation, inhalation, direct and indirect effects of oil ingestion
<i>Oceanodroma homochroa</i>	Ashy Storm-Petrel	High	yes	no	SC	yes	Oil effects on insulation, inhalation, direct and indirect effects of oil ingestion
<i>Oceanodroma melania</i>	Black Storm-Petrel	High	yes	no	no	no	Oil effects on insulation, inhalation, direct and indirect effects of oil ingestion
<i>Phaethon aethereus</i>	Red-billed Tropicbird	Low	no	no	no	no	Low probability of being oiled

<i>Phaethon rubricauda</i>	Red-tailed Tropicbird	Low	no	no		no	Low probability of being oiled
<i>Sula dactylatra</i>	Masked Booby	Low	no	no		no	Low probability of being oiled
<i>Sula leucogaster</i>	Brown Booby	Low	no	no		no	Low probability of being oiled
<i>Sula sula</i>	Red-footed Booby	Low	no	no		no	Low probability of being oiled
<i>Pelecanus occidentalis</i>	Brown Pelican	High	yes	no	D	no	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Pelecanus erythrorhynchos</i>	American White Pelican	Medium	yes	no		no	Moderate probability of being oiled
<i>Phalacrocorax penicillatus</i>	Brandt's Cormorant	High	yes	no		yes	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Phalacrocorax auritus</i>	Double-crested Cormorant	High	yes	no		yes	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Phalacrocorax pelagicus</i>	Pelagic Cormorant	High	yes	no		yes	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Fregata magnificens</i>	Magnificent Frigatebird	Low	no	no		no	Low probability of being oiled
<i>Fregata minor</i>	Great Frigatebird	Low	no	no		no	Low probability of being oiled
<i>Botaurus lentiginosus</i>	American Bittern	Medium	yes	no	SC	no	Moderate probability of being oiled
<i>Ardea herodias</i>	Great Blue Heron	Medium	yes	no		no	Moderate probability of being oiled
<i>Ardea alba</i>	Great Egret	Medium	yes	no		no	Moderate probability of being oiled
<i>Egretta thula</i>	Snowy Egret	Medium	yes	no		no	Moderate probability of being oiled
<i>Butorides virescens</i>	Green Heron	Medium	yes	no		no	Moderate probability of being oiled
<i>Nycticorax nycticorax</i>	Black-crowned Night-Heron	Medium	yes	no		no	Moderate probability of being oiled
<i>Cathartes aura</i>	Turkey Vulture	High	yes	no		no	Typical feather and ingestion effects
<i>Branta canadensis</i>	Canada Goose	Medium	yes	no	D	no	Typical feather and ingestion effects
<i>Branta bernicla</i>	Brant	High	yes	no		no	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Anas strepera</i>	Gadwall	Medium	yes	no		no	Moderate probability of being oiled
<i>Anas penelope</i>	Eurasian Wigeon	Medium	yes	no		no	Moderate probability of being oiled
<i>Anas americana</i>	American Wigeon	Medium	yes	no		no	Moderate probability of being oiled
<i>Anas platyrhynchos</i>	Mallard	Medium	yes	no		no	Moderate probability of being oiled
<i>Anas discors</i>	Blue-winged Teal	Medium	yes	no		no	Moderate probability of being oiled

<i>Anas cyanoptera</i>	Cinnamon Teal	Medium	yes	no		no	Moderate probability of being oiled
<i>Anas clypeata</i>	Northern Shoveler	Medium	yes	no		no	Moderate probability of being oiled
<i>Anas acuta</i>	Northern Pintail	Medium	yes	no		no	Moderate probability of being oiled
<i>Anas crecca</i>	Green-winged Teal	Medium	yes	no		no	Moderate probability of being oiled
<i>Aythya marila</i>	Greater Scaup	Medium	yes	no		no	Moderate probability of being oiled
<i>Aythya affinis</i>	Lesser Scaup	Medium	yes	no		no	Moderate probability of being oiled
<i>Histrionicus histrionicus</i>	Harlequin Duck	High	yes	no	SC	no	Typical feather and ingestion effects
<i>Melanitta perspicillata</i>	Surf Scoter	High	yes	no		no	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Melanitta fusca</i>	White-winged Scoter	High	yes	no		no	Typical feather and ingestion effects
<i>Melanitta nigra</i>	Black Scoter	High	yes	no		no	Typical feather and ingestion effects
<i>Clangula hyemalis</i>	Long-tailed Duck (Oldsquaw)	High	yes	no		no	Typical feather and ingestion effects
<i>Bucephala albeola</i>	Bufflehead	Medium	yes	no		no	Typical feather and ingestion effects
<i>Bucephala clangula</i>	Common Goldeneye	Medium	yes	no		no	Typical feather and ingestion effects
<i>Mergus serrator</i>	Red-breasted Merganser	High	yes	no		no	Typical feather and ingestion effects
	Common Merganser	Medium	yes	no		no	Moderate probability of being oiled
<i>Oxyura jamaicensis</i>	Ruddy Duck	Medium	yes	no		no	Moderate probability of being oiled
<i>Pandion haliaetus</i>	Osprey	High	yes	no		yes	Typical feather and ingestion effects
<i>Haliaeetus leucocephalus</i>	Bald Eagle	High	yes	no	T	yes	Typical feather and ingestion effects
	Golden Eagle	High	yes	no		yes	Typical feather and ingestion effects
<i>Circus cyaneus</i>	Northern Harrier	High	yes	no		yes	Typical feather and ingestion effects
<i>Falco columbarius</i>	Merlin	High	yes	no		yes	Typical feather and ingestion effects
<i>Falco peregrinus</i>	Peregrine Falcon	High	yes	no	D	yes	Typical feather and ingestion effects
<i>Laterallus jamaicensis</i>	Black Rail	Medium	yes	no	C	no	Moderate probability of being oiled
<i>Rallus limicola</i>	Virginia Rail	Medium	yes	no		no	Moderate probability of being oiled
<i>Coturnicops noveboracensis</i>	Yellow Rail	Medium	yes	no		no	Moderate probability of being oiled
<i>Porzana carolina</i>	Sora	Medium	yes	no		no	Moderate probability of being oiled
<i>Fulica americana</i>	American Coot	Medium	yes	no		no	Moderate probability of being oiled
<i>Pluvialis squatarola</i>	Black-bellied Plover	High	yes	no		no	Typical feather and ingestion effects
<i>Charadrius alexandrinus</i>	Snowy Plover	High	yes	no	T	yes	Typical feather and ingestion effects
<i>Charadrius semipalmatus</i>	Semipalmated Plover	High	yes	no		no	Typical feather and ingestion effects
<i>Charadrius vociferus</i>	Killdeer	Medium	yes	no		no	Moderate probability of being oiled
<i>Haematopus bachmani</i>	Black Oystercatcher	High	yes	no	SC	no	Typical feather and ingestion effects

<i>Recurvirostra americana</i>	American Avocet	High	yes	no	no	Typical feather and ingestion effects
<i>Tringa melanoleuca</i>	Greater Yellowlegs	High	yes	no	no	Typical feather and ingestion effects
<i>Catoptrophorus semipalmatus</i>	Willet	High	yes	no	no	Typical feather and ingestion effects
<i>Heteroscelus incanus</i>	Wandering Tattler	High	yes	no	no	Typical feather and ingestion effects
<i>Actitis macularia</i>	Spotted Sandpiper	High	yes	no	no	Typical feather and ingestion effects
<i>Numenius phaeopus</i>	Whimbrel	High	yes	no	SC	no Typical feather and ingestion effects
<i>Numenius americanus</i>	Long-billed Curlew	High	yes	no	SC	no Typical feather and ingestion effects
<i>Limosa fedoa</i>	Marbled Godwit	High	yes	no	SC	no Typical feather and ingestion effects
<i>Arenaria interpres</i>	Ruddy Turnstone	High	yes	no		no Typical feather and ingestion effects
<i>Arenaria melanocephala</i>	Black Turnstone	High	yes	no	SC	no Typical feather and ingestion effects
<i>Aphriza virgata</i>	Surfbird	High	yes	no		no Typical feather and ingestion effects
<i>Calidris canutus</i>	Red Knot	High	yes	no	SC	no Typical feather and ingestion effects
<i>Calidris alba</i>	Sanderling	High	yes	no		no Typical feather and ingestion effects
<i>Calidris mauri</i>	Western Sandpiper	High	yes	no		no Typical feather and ingestion effects
<i>Calidris minutilla</i>	Least Sandpiper	High	yes	no		no Typical feather and ingestion effects
<i>Calidris ptilocnemis</i>	Rock Sandpiper	High	yes	no		no Typical feather and ingestion effects
<i>Calidris alpina</i>	Dunlin	High	yes	no		no Typical feather and ingestion effects
<i>Limnodromus griseus</i>	Short-billed Dowitcher	High	yes	no		no Typical feather and ingestion effects
<i>Limnodromus scolopaceus</i>	Long-billed Dowitcher	High	yes	no		no Typical feather and ingestion effects
<i>Gallinago gallinago</i>	Common Snipe	High	yes	no		no Typical feather and ingestion effects
<i>Phalaropus lobatus</i>	Red-necked Phalarope	High	yes	no		no Typical feather and ingestion effects
	Wilson's Phalarope	High	yes	no		no Typical feather and ingestion effects
<i>Phalaropus fulicaria</i>	Red Phalarope	High	yes	no		no Typical feather and ingestion effects
<i>Catharacta maccormicki</i>	South Polar Skua	Low	no	no		no Low probability of being oiled
<i>Stercorarius pomarinus</i>	Pomarine Jaeger	Low	no	no		no Low probability of being oiled
<i>Stercorarius parasiticus</i>	Parasitic Jaeger	Low	no	no		no Low probability of being oiled
<i>Stercorarius longicaudus</i>	Long-tailed Jaeger	Low	no	no		no Low probability of being oiled
<i>Larus philadelphia</i>	Bonaparte's Gull	High	yes	no		no Typical feather and ingestion effects
<i>Larus heermanni</i>	Heermann's Gull	High	yes	no		no Typical feather and ingestion effects
<i>Larus canus</i>	Mew Gull	High	yes	no		no Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Larus delawarensis</i>	Ring-billed Gull	High	yes	no		no Typical feather and ingestion effects, experience shows frequent oiling

<i>Larus californicus</i>	California Gull	High	yes	no		yes	during oil spills Typical feather and ingestion effects
<i>Larus argentatus</i>	Herring Gull	High	yes	no		no	Typical feather and ingestion effects
<i>Larus thayeri</i>	Thayer's Gull	High	yes	no		no	Typical feather and ingestion effects
<i>Larus occidentalis</i>	Western Gull	High	yes	no		yes	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Larus glaucescens</i>	Glaucous-winged Gull	High	yes	no		no	Typical feather and ingestion effects
<i>Larus hyperboreus</i>	Glaucous Gull	High	yes	no		no	Typical feather and ingestion effects
<i>Xema sabini</i>	Sabine's Gull	High	yes	no		no	Typical feather and ingestion effects
<i>Creagrus furcatus</i>	Swallow-tailed Gull	High	yes	no		no	Typical feather and ingestion effects
<i>Rissa tridactyla</i>	Black-legged Kittiwake	High	yes	no		no	Typical feather and ingestion effects
<i>Sterna caspia</i>	Caspian Tern	High	yes	no		yes	Typical feather and ingestion effects
<i>Sterna elegans</i>	Elegant Tern	High	yes	no		yes	Typical feather and ingestion effects
<i>Sterna hirundo</i>	Common Tern	High	yes	no		no	Typical feather and ingestion effects
<i>Sterna paradisaea</i>	Arctic Tern	High	yes	no		no	Typical feather and ingestion effects
<i>Sterna forsteri</i>	Forster's Tern	High	yes	no		no	Typical feather and ingestion effects
<i>Sterna antillarum browni</i>	Least Tern	High	yes	no	E	yes	Typical feather and ingestion effects
<i>Uria aalge</i>	Common Murre	High	yes	no		yes	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Uria lomvia</i>	Thick-billed Murre	High	yes	no		no	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Cephus columba</i>	Pigeon Guillemot	High	yes	no		yes	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Brachyramphus marmoratus</i>	Marbled Murrelet	High	yes	no	T	no	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Brachyramphus perdix</i>	Long-billed Murrelet	High	yes	no		no	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Synthliboramphus hypoleucus</i>	Xantus's Murrelet	High	yes	no	SC	no	Typical feather and ingestion effects, experience shows frequent oiling during oil spills

<i>Synthliboramphus craveri</i>	Craveri's Murrelet	High	yes	no		no	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Synthliboramphus antiquus</i>	Ancient Murrelet	High	yes	no		no	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Ptychoramphus aleuticus</i>	Cassin's Auklet	High	yes	no	SC	yes	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Aethia psittacula</i>	Parakeet Auklet	High	yes	no		no	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Aethia pusilla</i>	Least Auklet	High	yes	no		no	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Aethia cristatella</i>	Crested Auklet	High	yes	no		no	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Cerorhinca monocerata</i>	Rhinoceros Auklet	High	yes	no		yes	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Fratercula corniculata</i>	Horned Puffin	High	yes	no		no	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Fratercula cirrhata</i>	Tufted Puffin	High	yes	no		yes	Typical feather and ingestion effects, experience shows frequent oiling during oil spills
<i>Asio flammeus</i>	Short-eared Owl	Low	no	no		no	Low probability of being oiled
<i>Ceryle alcyon</i>	Belted Kingfisher	Medium	yes	no		yes	Moderate probability of being oiled
<i>Sayornis nigricans</i>	Black Phoebe	Low	no	no		no	Low probability of being oiled
<i>Sayornis saya</i>	Say's Phoebe	Low	no	no		no	Low probability of being oiled
<i>Corvus corax</i>	Common Raven	Medium	yes	no		yes	Typical feather and ingestion effects
<i>Corvus brachyrhynchos</i>	Common crow	Low	yes	no		yes	Low probability of being oiled
<i>Eremophila alpestris</i>	Horned Lark	Low	no	no		no	Low probability of being oiled
<i>Tachycineta bicolor</i>	Tree Swallow	Low	no	no		no	Low probability of being oiled
<i>Stelgidopteryx serripennis</i>	Northern Rough-winged Swallow	Low	no	no		no	Low probability of being oiled

<i>Petrochelidon pyrrhonota</i>	Cliff Swallow	Low	no	no	no	Low probability of being oiled
<i>Hirundo rustica</i>	Barn Swallow	Low	no	no	no	Low probability of being oiled
<i>Salpinctes obsoletus</i>	Rock Wren	Low	no	no	no	Low probability of being oiled
<i>Cistothorus palustris</i>	Marsh Wren	Low	no	no	no	Low probability of being oiled
<i>Anthus rubescens</i>	American Pipit	Low	no	no	no	Low probability of being oiled
<i>Dendroica coronata</i>	Yellow-rumped Warbler	Low	no	no	no	Low probability of being oiled
<i>Passerculus sandwichensis</i>	Savannah Sparrow	Low	no	no	no	Low probability of being oiled
<i>Melospiza melodia</i>	Song Sparrow	Low	no	no	no	Low probability of being oiled
<i>Melospiza georgiana</i>	Swamp Sparrow	Low	no	no	no	Low probability of being oiled
<i>Agelaius phoeniceus</i>	Red-winged Blackbird	Low	no	no	no	Low probability of being oiled
<i>Sturnella neglecta</i>	Western Meadowlark	Low	no	no	no	Low probability of being oiled

---

**Fish**

Scientific Name	Common Name	Sensitivity to oil	More Benefit from Dispersant	More Harm from Dispersant	ESA listing	Breeds locally	Reason
<i>Eptatretus deani</i>	Black Hagfish						
<i>Eptatretus stoutii</i>	Pacific Hagfish						
<i>Lampreta tridentata</i>	Pacific Lamprey	Medium	yes	no	SC	yes	Effects of smothering of oil inside estuaries on eggs and larvae
<i>Lampetra ayersii</i>	Western River Lamprey						
<i>Hydrolagus colliei</i>	Spotted Ratfish						
<i>Hexanchus griseus</i>	Bluntnose Sixgill Shark						
<i>Notorynchus cepedianus</i>	Broadnose Sevengill Shark						
<i>Echinorhinus cookei</i>	Prickly Shark						
<i>Squalus acanthias</i>	Spiny Dogfish						
<i>Somniosus pacificus</i>	Pacific Sleeper Shark						
<i>Squatina californica</i>	Pacific Angel Shark						
<i>Alopias vulpinus</i>	Thresher Shark						
<i>Cetorhinus maximus</i>	Basking Shark						
<i>Carcharodon carcharias</i>	White Shark	Low	yes	no		yes	Limited potential exposure to oil
<i>Isurus oxyrinchus</i>	Shortfin Mako						
<i>Lamna ditropis</i>	Salmon Shark						
<i>Apristurus brunneus</i>	Brown Catshark						

<i>Apristurus kampae</i>	Longnose Catshark							
<i>Parmaturus xaniurus</i>	Filetail Catshark							
<i>Galeorhinus galeus</i>	Tope or Soupfin Shark							
<i>Mustelus californicus</i>	Gray Smoothhound							
<i>Mustelus henlei</i>	Brown Smoothhound							
<i>Triakis semifasciata</i>	Leopard Shark							
<i>Prionace glauca</i>	Blue Shark							
<i>Torpedo californica</i>	Pacific Electric Ray							
<i>Rhinobatos productus</i>	Shovelnose Guitarfish							
<i>Platyrhinoidis triseriata</i>	Pacific Thornback							
<i>Amblyraja badia</i>	Broad skate							
<i>Bathyraja abyssicola</i>	Deepsea Skate							
<i>Bathyraja interrupta</i>	Sandpaper Skate							
<i>Bathyraja spinosissima</i>	White Skate							
<i>Bathyraja trachura</i>	Black Skate							
<i>Raja binoculata</i>	Big Skate							
<i>Raja inornata</i>	California Skate							
<i>Raja rhina</i>	Longnose Skate							
<i>Raja stellulata</i>	Starry Skate							
<i>Dasyatis diptera</i>	Diamond Stingray							
<i>Dasyatis violacea</i>	Pelagic Stingray							
<i>Urolophus halleri</i>	Round Stingray							
<i>Myliobatis californica</i>	Bat Ray							
<i>Acipenser medirostris</i>	Green Sturgeon	Low	yes	no	T	yes	Bottom dweller, highly mobile, juveniles of limited concern	
<i>Acipenser transmontanus</i>	White Sturgeon						Vulnerability of egg and larvae in the upper water column to dissolved fraction of oil	
<i>Albula vulpes</i>	Bonefish							
<i>Ophichthus triserialis</i>	Pacific Snake Eel							
<i>Ophichthus zaphochir</i>	Yellow Snake Eel							
<i>Nemichthys scolopaceus</i>	Slender Snipe Eel							
<i>Serrivomer sector</i>	Sawtooth Snipe Eel							
<i>Cyema atrum</i>	Bobtail Snipe Eel							

<i>Engraulis mordax</i>	Northern Anchovy							
<i>Alosa sapidissima</i>	American Shad							
<i>Clupea pallasii</i>	Pacific Herring	High	yes	no		yes	Toxic effects of oil on eggs at the intertidal	
<i>Sardinops sagax</i>	Pacific Sardine							
<i>Argentina sialis</i>	Pacific Argentine							
<i>Bathylagoides wesethi</i>	Snubnose Blacksmelt							
<i>Bathylagus pacificus</i>	Pacific Blacksmelt							
<i>Leuroglossus stilbius</i>	California Smoothtongue							
<i>Lipolagus ochotensis</i>	Popeye Blacksmelt							
<i>Pseudobathylagus milleri</i>	Robust Blacksmelt							
<i>Macropinna microstoma</i>	Pacific Barreleye							
<i>Alepocephalus tenebrosus</i>	California Slickhead							
<i>Talismania bifurcata</i>	Threadfin Slickhead							
<i>Sagamichthys abei</i>	Shining Tubeshoulder							
<i>Allosmerus elongatus</i>	Whitebait Smelt							
<i>Hypomesus pretiosus</i>	Surf Smelt							
<i>Spirinchus starksi</i>	Night Smelt							
<i>Spirinchus thaleichthys</i>	Longfin Smelt	?						
<i>Thaleichthys pacificus</i>	Eulachon	Low	yes	no	T	no	Rare species in central California	
<i>Oncorhynchus gorbuscha</i>	Pink Salmon							
<i>Oncorhynchus keta</i>	Chum Salmon	Low	yes	no	T	no	Rare species in central California	
<i>Oncorhynchus kisutch</i>	Coho Salmon [Silver Salmon]	Medium	yes	no	E&T	yes	Potential effect of oil on smolts using local lagoons	
<i>Oncorhynchus mykiss</i>	Rainbow Trout [Steelhead]	Medium	yes	no	E&T	yes	Potential effect of oil on smolts using local lagoons	
<i>Oncorhynchus nerka</i>	Sockeye Salmon							
<i>Oncorhynchus tshawytscha</i>	Chinook Salmon	Low	yes	no	E&T	yes	Potential effect of oil on smolts using local lagoons	
<i>Cyclothone acclinidens</i>	Benttooth Bristlemouth							
<i>Cyclothone signata</i>	Showy Bristlemouth							
<i>Daphnos oculatus</i>	Bigeye Lightfish							
<i>Argyropelecus affinis</i>	Slender Hatchetfish							
<i>Argyropelecus hemigymnus</i>	Spurred Hatchetfish							

<i>Argyropelecus lychnus</i>	Silver Hatchetfish
<i>Argyropelecus sladeni</i>	Silvery Hatchetfish
<i>Sternoptyx spp.</i>	Dollar Hatchetfishes
<i>Aristostomias scintillans</i>	Shiny Loosejaw
<i>Bathophilus flemingi</i>	Highfin Dragonfish
<i>Chauliodus macouni</i>	Pacific Viperfish
<i>Idiacanthus antrostomas</i>	Pacific Blackdragon
<i>Tactostoma macropus</i>	Longfin Dragonfish
<i>Benthalbella dentata</i>	Northern Pearleye
<i>Synodus lucioceps</i>	California Lizardfish
<i>Lestidium ringens</i>	Slender Barricudina
<i>Anopterus pharao</i>	Daggertooth
<i>Alepisaurus ferox</i>	Longnose Lancetfish
<i>Ceratoscopelus townsendi</i>	Dogtooth Lampfish
<i>Diaphus theta</i>	California Headlightfish
<i>Diogenes laternatus</i>	Diogenes Lanternfish
<i>Nannobranchium regale</i>	Pinpoint Lampfish
<i>Nannobranchium ritteri</i>	Broadfin Lampfish
<i>Notoscopelus resplendens</i>	Patchwork Lampfish
<i>Protomyctophum crockeri</i>	California Flashlightfish
<i>Protomyctophum thompsoni</i>	Bigeye Lanternfish
<i>Stenobranchius leucopsaurus</i>	Northern Lampfish
<i>Tarletonbaenia crenularis</i>	Blue Lanternfish
<i>Triphoturus mexicanus</i>	Mexican Lampfish
<i>Lampris regius</i>	Opah
<i>Desmodema lorum</i>	Whiptail Ribbonfish
<i>Trachipterus altivelis</i>	King-of-the-salmon
<i>Chilara taylori</i>	Spotted Cusk Eel
<i>Brosmophycis marginata</i>	Red Brotula
<i>Albatrossia pectoralis</i>	Giant Grenadier
<i>Coelorinchus scaphopsis</i>	Shoulderspot Grenadier
<i>Coryphaenoides acrolepis</i>	Pacific Grenadier
<i>Nezumia stelgidolepis</i>	California Grenadier

<i>Antimora microlepis</i>	Finescale Codling
<i>Physiculus rastrelliger</i>	Hundred-Fathom Codling
<i>Merluccius productus</i>	Pacific Hake
<i>Gadus microcephalus</i>	Pacific Cod
<i>Microgadus proximus</i>	Pacific Tomcod
<i>Theragra chalcogramma</i>	Walleye Pollock
<i>Porichthys notatus</i>	Plainfin Midshipman
<i>Atherinops affinis</i>	Topsmelt
<i>Atherinopsis californiensis</i>	Jacksmelt
<i>Leuresthes tenuis</i>	California Grunion
<i>Strongylura exilis</i>	California Needlefish
<i>Cololabis saira</i>	Pacific Saury
<i>Cheilopogon pinnatibarbatu</i>	Smallhead Flyingfish
<i>Melamphaes lugubris</i>	Highsnout Bigscale
<i>Poromitra crassiceps</i>	Crested Bigscale
<i>Scopeloberyx robustus</i>	Longjaw Bigscale
<i>Scopelogadus mizolepis</i>	Twospine Bigscale
<i>Anoplogaster cornuta</i>	Fangtooth
<i>Alloctytus folletti</i>	Oxeye oreo
<i>Aulorhynchus flavidus</i>	Tubesnout
<i>Gasterosteus aculeatus</i>	Threespine Stickleback
<i>Cosmocampus arctus</i>	Snubnose Pipefish
<i>Syngnathus californiensis</i>	Kelp Pipefish
<i>Syngnathus leptorhynchus</i>	Bay Pipefish
<i>Sebastes aleutianus</i>	Rougheye Rockfish
<i>Sebastes alutus</i>	Pacific Ocean Perch
<i>Sebastes atrovirens</i>	Kelp Rockfish
<i>Sebastes auriculatus</i>	Brown Rockfish
<i>Sebastes aurora</i>	Aurora Rockfish
<i>Sebastes babcocki</i>	Redbanded Rockfish
<i>Sebastes brevispinis</i>	Silvergray Rockfish
<i>Sebastes carnatus</i>	Gopher Rockfish
<i>Sebastes caurinus</i>	Copper Rockfish

<i>Sebastes chlorostictus</i>	Greenspotted rockfish						
<i>Sebastes chrysomelas</i>	Black-and-Yellow Rockfish						
<i>Sebastes constellatus</i>	Starry Rockfish						
<i>Sebastes crameri</i>	Darkblotched Rockfish						
<i>Sebastes dallii</i>	Calico Rockfish						
<i>Sebastes diploproa</i>	Splitnose Rockfish						
<i>Sebastes elongatus</i>	Greenstriped Rockfish						
<i>Sebastes ensifer</i>	Swordspine Rockfish						
<i>Sebastes entomelas</i>	Widow Rockfish						
<i>Sebastes eos</i>	Pink Rockfish						
<i>Sebastes flavidus</i>	Yellowtail rockfish						
<i>Sebastes goodei</i>	Chilipepper						
<i>Sebastes helvomaculatus</i>	Rosethorn Rockfish						
<i>Sebastes hopkinsi</i>	Squarespot Rockfish						
<i>Sebastes jordani</i>	Shortbelly Rockfish						
<i>Sebastes levis</i>	Cowcod	Low	yes	no		yes	Deep water species with limited potential exposure to oil
<i>Sebastes maliger</i>	Quillback Rockfish						
<i>Sebastes melanops</i>	Black Rockfish						
<i>Sebastes melanostomus</i>	Blackgill Rockfish						
<i>Sebastes miniatus</i>	Vermilion Rockfish						
<i>Sebastes mystinus</i>	Blue Rockfish						
<i>Sebastes nebulosus</i>	China Rockfish						
<i>Sebastes nigrocinctus</i>	Tiger Rockfish						
<i>Sebastes ovalis</i>	Speckled Rockfish						
<i>Sebastes paucispinis</i>	Bocaccio	Low	yes	no	SC	yes	Deep water species with limited potential exposure to oil
<i>Sebastes phillipsi</i>	Chameleon Rockfish						
<i>Sebastes pinniger</i>	Canary Rockfish	Low	yes	no		yes	Deep water species with limited potential exposure to oil
<i>Sebastes proriger</i>	Redstripe Rockfish						
<i>Sebastes rastrelliger</i>	Grass Rockfish						
<i>Sebastes rosaceus</i>	Rosy Rockfish						
<i>Sebastes rosenblatti</i>	Greenblotched Rockfish						

<i>Sebastes ruberrimus</i>	Yelloweye Rockfish	Low	yes	no	yes	Deep water species with limited potential exposure to oil
<i>Sebastes rubrivinctus</i>	Flag Rockfish					
<i>Sebastes rufus</i>	Bank Rockfish					
<i>Sebastes saxicola</i>	Stripetail Rockfish					
<i>Sebastes semicinctus</i>	Halfbanded Rockfish					
<i>Sebastes serranoides</i>	Olive Rockfish					
<i>Sebastes serriceps</i>	Treefish					
<i>Sebastes wilsoni</i>	Pygmy Rockfish					
<i>Sebastes zacentrus</i>	Sharpchin Rockfish					
<i>Sebastolobus alascanus</i>	Shortspine Thornyhead					
<i>Sebastolobus altivelis</i>	Longspine Thornyhead					
<i>Prionotus stephanophrys</i>	Lumptail Searobin					
<i>Anoplopoma fimbria</i>	Sablefish					
<i>Erilepis zonifer</i>	Skilfish					
<i>Hexagrammos decagrammus</i>	Kelp Greenling					
<i>Hexagrammos superciliosus</i>	Rock Greenling					
<i>Ophiodon elongaus</i>	Lingcod	Low	yes	no	yes	Deep water species with limited potential exposure to oil
<i>Oxylebius pictus</i>	Painted Greenling					
<i>Zaniolepis frenata</i>	Shortspine Combfish					
<i>Zaniolepis latipinnis</i>	Longspine Combfish					
<i>Rhamphocottus richardsonii</i>	Grunt Sculpin					
<i>Artedius corallinus</i>	Corraline Sculpin					
<i>Artedius fenestralis</i>	Padded Sculpin					
<i>Artedius harringtoni</i>	Scalyhead Sculpin					
<i>Artedius lateralis</i>	Smoothhead Sculpin					
<i>Artedius notospilotus</i>	Bonyhead Sculpin					
<i>Ascelichthys rhodorus</i>	Rosylip Sculpin					
<i>Chitonotus pugetensis</i>	Roughback Sculpin					
<i>Clinocottus acuticeps</i>	Sharpnose Sculpin					
<i>Clinocottus analis</i>	Wooly Sculpin					
<i>Clinocottus embryum</i>	Calico Sculpin					
<i>Clinocottus globiceps</i>	Mosshead Sculpin					

<i>Clinocottus recalvus</i>	Bald Sculpin
<i>Enophrys bison</i>	Buffalo Sculpin
<i>Enophrys taurina</i>	Bull Sculpin
<i>Hemilepidotus hemilepidotus</i>	Red Irishlord
<i>Hemilepidotus spinosus</i>	Brown Irishlord
<i>Icelinus burchami</i>	Dusky Sculpin
<i>Icelinus filamentosus</i>	Threadfin Sculpin
<i>Icelinus oculatus</i>	Frogmouth Sculpin
<i>Icelinus quadriseriatus</i>	Yellowchin Sculpin
<i>Icelinus tenuis</i>	Spotfin Sculpin
<i>Jordania zonope</i>	Longfin Sculpin
<i>Leptocottus armatus</i>	Staghorn Sculpin
<i>Oligocottus maculosus</i>	Tidepool Sculpin
<i>Oligocottus rimensis</i>	Saddleback Sculpin
<i>Oligocottus rubellio</i>	Rosy Sculpin
<i>Oligocottus snyderi</i>	Fluffy Sculpin
<i>Orthonopias triacis</i>	Snubnose Sculpin
<i>Paricelinus hopliticus</i>	Thornback Sculpin
<i>Radulinus boleoides</i>	Darter Sculpin
<i>Scorpaenichthys marmoratus</i>	Cabazon Sculpin
<i>Synchirus gilli</i>	Manacled Sculpin
<i>Belpsius cirrhosus</i>	Silverspotted Sculpin
<i>Nautichthys oculoasciatus</i>	Sailfin Sculpin
<i>Agonopsis vulsa</i>	Northern Spearnose Poacher
<i>BathYGONUS pentacanthus</i>	Bigeye Poacher
<i>BothrAGONUS swanii</i>	Rockhead
<i>Chesnonia verrucosa</i>	Warty Poacher
<i>HypsAGONUS mozinoi</i>	Kelp Poacher
<i>OdontOPYXIS trispinosa</i>	Pygmy Poacher
<i>Pallasina barbata</i>	Tube-nose Poacher
<i>Stellerina xyosterna</i>	Pricklebreast Poacher
<i>Xeneretmus latifrons</i>	Blackedge Poacher
<i>Xeneretmus leiops</i>	Smootheye Poacher

<i>Xeneretmus triacanthus</i>	Bluespotted Poacher
<i>Psychrolutes phrictus</i>	Blob Sculpin
<i>Careproctus melanurus</i>	Blacktail Snailfish
<i>Liparis florum</i>	Tidepool Snailfish
<i>Liparis fuscensis</i>	Slipskin Snailfish
<i>Liparis mucosus</i>	Slimy Snailfish
<i>Liparis adiastolus</i>	Southern Ringtail Snailfish
<i>Liparis pulchellus</i>	Showy Snailfish
<i>Morone saxatilis</i>	Striped Bass
<i>Stereolepis gigas</i>	Giant Sea Bass
<i>Mycteroperca xenarcha</i>	Broomtail Grouper
<i>Paralabrax clathratus</i>	Kelp Bass
<i>Caulotilus princeps</i>	Ocean Whitefish
<i>Remora albescens</i>	White Suckerfish
<i>Remora australis</i>	Whalesucker
<i>Remora remora</i>	Remora
<i>Coryphaena hippurus</i>	Dolphinfish
<i>Trachurus symmetricus</i>	Jack Mackerel
<i>Seriola lalandi</i>	Yellowtail Jack
<i>Brama japonica</i>	Pacific Pomfret
<i>Caristius macropus</i>	Veilfin
<i>Atractoscion nobilis</i>	White Seabass
<i>Genyonemus lineatus</i>	White Croaker
<i>Seriphus politus</i>	Queenfish
<i>Pseudopentaceros wheeleri</i>	North Pacific Armorhead
<i>Girella nigricans</i>	Opaleye
<i>Medialuna californiensis</i>	Halfmoon
<i>Amphistichus argenteus</i>	Barred Surfperch
<i>Amphistichus koelzi</i>	Calico Surfperch
<i>Amphistichus rhodotus</i>	Redtail Surfperch
<i>Brachyistius frenatus</i>	Kelp Perch
<i>Cymatogaster aggregata</i>	Shiner Perch
<i>Damalichthys vacca</i>	Pile Perch

<i>Embiotoca jacksoni</i>	Black Perch						
<i>Embiotoca lateralis</i>	Striped Seaperch						
<i>Hyperprosopon anale</i>	Spotfin Surfperch						
<i>Hyperprosopon argenteum</i>	Walleye Surfperch						
<i>Hyperprosopon ellipticum</i>	Silver Surfperch						
<i>Hypsurus caryi</i>	Rainbow Seaperch						
<i>Micrometrus aurora</i>	Reef Perch						
<i>Micrometrus minimus</i>	Dwarf Perch						
<i>Phanerodon atripes</i>	Sharpnose Seaperch						
<i>Phanerodon furcatus</i>	White Seaperch						
<i>Rhacochilus toxotes</i>	Rubberlip Seaperch						
<i>Zalemnius rosaceus</i>	Pink Seaperch						
<i>Oxyjulis californica</i>	Señorita						
<i>Semicossyphus pulcher</i>	California Sheephead						
<i>Rathbunella alleni</i>	Stripefin Ronquil						
<i>Ronquilus jordani</i>	Northern Ronquil						
<i>Bothrocara brunneum</i>	Twoline Eelpout						
<i>Bothrocara molle</i>	Soft Eelpout						
<i>Embryx crotalina</i>	Flatcheek Eelpout						
<i>Lycodapus fierasfer</i>	Blackmouth Eelpout						
<i>Lycodapus mandibularis</i>	Pallid Eelpout						
<i>Lycodes cortezianus</i>	Bigfin Eelpout						
<i>Lycodes diapterus</i>	Black Eelpout						
<i>Lycodopsis pacifica</i>	Blackbelly Eelpout						
<i>Lyonema barbatus</i>	Bearded Eelpout						
<i>Melanostigma pammelas</i>	Midwater Eelpout						
<i>Anoplarchus purpureus</i>	High Cockscomb						
<i>Cebidichthys violaceus</i>	Monkeyface Prickleback	High	yes	no	yes	Intertidal and subtidal species likely to be affected by toxicity of oil	
<i>Chirolophis nugator</i>	Mosshead Warbonnet						
<i>Kasatkia seigeli</i>	Sixspot Prickleback						
<i>Phytichthys chirus</i>	Ribbon Prickleback						
<i>Plectrobranchius evides</i>	Bluebarred Prickleback						
<i>Poroclinus rothrocki</i>	Whitebarred Prickleback						

<i>Xiphister atropurpureus</i>	Black Prickleback							
<i>Xiphister mucosus</i>	Rock Prickleback							
<i>Apodichthys flavidus</i>	Penpoint Gunnel							
<i>Apodichthys fucorum</i>	Rockweed Gunnel							
<i>Pholis ornata</i>	Saddleback Gunnel							
<i>Pholis schultzi</i>	Red Gunnel							
<i>Anarrhichthys ocellatus</i>	Wolf-Eel							
<i>Zaprora silenus</i>	Prowfish							
<i>Scytalina cerdale</i>	Graveldiver							
<i>Trichodon trichodon</i>	Pacific Sandfish							
<i>Ammodytes hexapterus</i>	Pacific Sand Lance							
<i>Gibbonsia metzi</i>	Striped Kelpfish							
<i>Gibbonsia montereyensis</i>	Crevice Kelpfish							
<i>Heterostichus rostratus</i>	Giant Kelpfish							
<i>Neoclinus blanchardi</i>	Sarcastic Fringehead							
<i>Neoclinus uniornatus</i>	Onespot Fringehead							
<i>Icosteus aenigmaticus</i>	Ragfish							
<i>Gobiesox meandricus</i>	Northern Clingfish							
<i>Rimicola muscarum</i>	Kelp Clingfish							
<i>Acanthogobius flavimanus</i>	Yellowfin Goby							
<i>Clevelandia ios</i>	Arrow Goby							
<i>Eucyclogobius newberryi</i>	Tidewater Goby	High	yes	no	E	yes	Toxic effects of oil, smothering of tide pools by oil	
<i>Gillichthys mirabilis</i>	Longjaw Mudsucker							
<i>Ilypnus gilberti</i>	Cheekspot Goby							
<i>Lepidogobius lepidus</i>	Bay Goby							
<i>Coryphopterus nicholsii</i>	Blackeye Goby							
<i>Luvarus imperialis</i>	Louvar							
<i>Sphyræna argentea</i>	Pacific Barracuda							
<i>Lepidocybrium flavobrunneum</i>	Escolar							
<i>Lepidopus fitchi</i>	Pacific Scabbardfish							
<i>Katsuwonus pelamis</i>	Skipjack Tuna							
<i>Sarda chiliensis</i>	Pacific Bonito							
<i>Scomber japonicus</i>	Pacific Chub Mackerel							

<i>Thunnus alalunga</i>	Albacore
<i>Thunnus obesus</i>	Bigeye Tuna
<i>Thunnus orientalis</i>	Pacific Bluefin Tuna
<i>Xiphias gladius</i>	Swordfish
<i>Tetrapturus angustirostris</i>	Shortbill Spearfish
<i>Tetrapturus audax</i>	Striped Marlin
<i>Icichthys lockingtoni</i>	Medusafish
<i>Tetrogonurus cuvieri</i>	Smalleye Squaretail
<i>Peprilus simillimus</i>	Pacific Pompano
<i>Citharichthys sordidus</i>	Pacific Sanddab
<i>Citharichthys stigmaeus</i>	Speckled Sanddab
<i>Paralichthys californicus</i>	California Halibut
<i>Atheresthes stomias</i>	Arrowtooth Flounder
<i>Clidoderma asperrimum</i>	Roughscale Sole
<i>Embassichthys bathybius</i>	Deepsea Sole
<i>Eopsetta jordani</i>	Petrale Sole
<i>Glyptocephalus zachirus</i>	Rex Sole
<i>Hippoglossoides elassodon</i>	Flathead Sole
<i>Hippoglossus stenolepis</i>	Pacific Halibut
<i>Isopsetta isolepis</i>	Butter Sole
<i>Lepidopsetta bilineata</i>	Rock Sole
<i>Lyopsetta exilis</i>	Slender Sole
<i>Microstomus pacificus</i>	Dover Sole
<i>Parophrys vetulus</i>	English Sole
<i>Platichthys stellatus</i>	Starry Flounder
<i>Pleuronichthys coenosus</i>	C-O Sole
<i>Pleuronichthys decurrens</i>	Curlfin Sole
<i>Pleuronichthys guttulatus</i>	Diamond Turbot
<i>Pleuronichthys verticalis</i>	Hornyhead Turbot
<i>Psettichthys melanostictus</i>	Sand Sole
<i>Reinhardtius hippoglossoides</i>	Greenland Halibut
<i>Symphurus atricauda</i>	California Tonguefish
<i>Balistes polylepis</i>	Finescale Triggerfish

<i>Lagocephalus lagocephalus</i>	Oceanic Pufferfish
<i>Diodon holocanthus</i>	Balloonfish
<i>Mola mola</i>	Ocean Sunfish

---

**Reptiles**

Scientific Name	Common Name	Sensitivity to oil	More Benefit from Dispersant	More Harm from Dispersant	ESA listing	Breeds locally	Reason
<i>Chelonia mydas</i>	Green Sea Turtle	Low	yes	no	T	no	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Lepidochelys olivacea</i>	Pacific (Olive) Ridley	Low	yes	no	T	no	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Caretta caretta</i>	Loggerhead Turtle	Low	yes	no	T	no	Potentially affected by oil fumes, direct and indirect effects of oil ingestion
<i>Dermochelys coriacea</i>	Leatherback Turtle	Low	yes	no	E	no	Potentially affected by oil fumes, direct and indirect effects of oil ingestion

**INVERTEBRATES**

	Sensitivity to oil	More Benefit from Dispersant	More Harm from Dispersant	ESA listing	Breeds locally	Reason
Note: All species below that are included in zooplankton and ichthyoplankton	Low	yes	no		yes	Localized impacts and rapid repopulation expected

**Annelida**

Scientific Name	Common Name	Sensitivity to oil	More Benefit from Dispersant	More Harm from Dispersant	ESA listing	Breeds locally	Reason
<i>Arabella iricolor</i> <i>Cheilonereis cyclurus</i> <i>Errantia spp.</i> <i>Nereis guberi</i> <i>Phragmatopoma californica</i>	Polycheate						

*Phyllochaetopterus prolifica*  
*Platynereis bicanaliculata*  
*Serpula vermicularis* Tube worm  
*Spirorbis borealis*  
*Stylanthea prophyra*  
*Terribellidae*  
*Thelepus crispus*  
*Typosyllis aciculata*

---

**Arthropoda**

---

Scientific Name	Common Name	Sensitivity to oil	More Benefit from Dispersant	More Harm from Dispersant	ESA listing	Breeds locally	Reason
<i>Acanthomysis sp.</i>							
<i>Achelia chelata</i>							
<i>Achelia nudiscula</i>							
<i>Achelia spinoseta</i>							
<i>Allorchestes anceps</i>							
<i>Alpheus dentipes</i>							
<i>Ammothea hilgendorfi</i>							
<i>Amphiodia occidentalis</i>							
<i>Amphissa columbiana</i>							
<i>Amphissa versicolor</i>							
<i>Anatanais normani</i>							
<i>Balanus amphitrite</i>							
<i>Balanus cariosus</i>	Barnacle						
<i>Balanus glandula</i>	Barnacle						
<i>Balanus nubilus</i>	Barnacle						
<i>Balanus sp.</i>							
<i>Cancer antennarius</i>							
<i>Cancer magister</i>		Low	yes	no		yes	Wide distribution, bottom dweller, not likely to encounter oil
<i>Cancer productus</i>		Moderate	yes	no		yes	Wide distribution, shallow water, maybe affected by oil on intertidal and

subtidal

*Caprella californica*  
*Chthamalus dalli*  
*Cirolana harfordi*  
*Elasmopus serricatus*  
*Euphausia pacifica* Krill  
*Exosphaeroma inornata*  
*Exosphaeroma rhomburum*  
*Fabia subquadrata*  
*Hemigrapsus nudus*  
*Hyale frequens*  
*Hyale grandicornis*  
*Ianiropsis kincaidi*  
*Idotea fewkesi*  
*Idotea resecata*  
*Idotea schmitti*  
*Idotea sp.*  
*Idotea stenops*  
*Idotea urotoma*  
*Idotea wosnesenskii*  
*Lecythorhynchus hilgendorfi*  
*Ligia occidentalis*  
*Ligia pallasii*  
*Limnoria algarum*  
*Littorophiloscia richardsonae*  
*Lophopanopeus leucomanus*  
*Loxorhynchus crispatus* Crab  
*Melita californica*  
*Metacaprella anomala*  
*Metacaprella kennerlyi*  
*Nebalia kensleyi*  
*Nymphopsis spinosissima*  
*Oedignathus inermis*

<i>Oligochinus lighti</i>	
<i>Pachycheles rudis</i>	
<i>Pachygrapsus crassipes</i>	Crab
<i>Pachygrapsus nudus</i>	
<i>Pagurus granosimanus</i>	
<i>Pagurus hirsutiusculus</i>	Hermit crab
<i>Pagurus samuelensis</i>	
<i>Pagurus sp.</i>	
<i>Paracerceis cordata</i>	
<i>Paradynoides benedicti</i>	
<i>Parallorchestes ochotensis</i>	
<i>Paranthura elegans</i>	
<i>Paraxanthia taylorii</i>	
<i>Petrolisthes cinctipes</i>	
<i>Pinnixa franciscana</i>	
<i>Pollicipes polymerus</i>	
<i>Polycheria osborni</i>	
<i>Porcellio americanus</i>	
<i>Pugetia fragilissima</i>	Crab
<i>Pugettia gracilis</i>	Crab
<i>Pugettia producta</i>	Crab
<i>Pycnogonum rickettsi</i>	Sea spider
<i>Pycnogonum stearnsi</i>	Sea spider
<i>Scyra acutifrons</i>	Crab
<i>Semibalanus cariosus</i>	Barnacle
<i>Semibalanus sp.</i>	
<i>Tetraclita rubescens</i>	Barnacle
<i>Thysanoessa spinifera</i>	Krill

---

**Chordata**

Scientific Name	Common Name	Sensitivity to oil	More Benefit from Dispersant	More Harm from Dispersant	ESA listing	Breeds locally	Reason
-----------------	-------------	--------------------	------------------------------	---------------------------	-------------	----------------	--------

---

<i>Aplidium arenatum</i>	
<i>Aplidium californicum</i>	Tunicate
<i>Cystodytes lobatus</i>	Tunicate
<i>Didemnum carnulentum</i>	Tunicate
<i>Polyclinum planum</i>	
<i>Pycnoclayella stanleyi</i>	Tunicate
<i>Ritterella aequalisphonis</i>	Tunicate

---

**Cnidaria**

Scientific Name	Common Name	Sensitivity to oil	More Benefit from Dispersant	More Harm from Dispersant	ESA listing	Breeds locally	Reason
<i>Abietinaria sp.</i>	Fern hydroid						
<i>Aglaophenia inconspicua</i>							
<i>Aglaophenia latirostris</i>	Ostrich-plume hydroid						
<i>Aglaophenia sp.</i>							
<i>Anthopleura elegantissima</i>	Aggregating anemone						
<i>Anthopleura xanthogrammica</i>	Giant green anemone						
<i>Aurelia aurita</i>							
<i>Balanophyllia elegans</i>	Orange cup coral						
<i>Corynactis californica</i>							
<i>Epiactis prolifera</i>	Poliferating anemone						
<i>Eudendrium californicum</i>							
<i>Garveia annulata</i>							
<i>Metridium senile</i>	White-plumed anemone						
<i>Obelia sp.</i>							
<i>Sertularella turgida</i>							
<i>Sertularia sp.</i>							
<i>Stylatula elongata</i>	Sea pen						
<i>Tealia crassicornis</i>							
<i>Tealia lofotensis</i>							
<i>Tubularia crocea</i>							
<i>Urticina crassicornia</i>							

*Urticina lofotensis*

---

**Echinodermata**

---

Scientific Name	Common Name	Sensitivity to oil	More Benefit from Dispersant	More Harm from Dispersant	ESA listing	Breeds locally	Reason
<i>Amphipholis squamata</i>							
<i>Asterina miniata</i>							
<i>Cucumaria curata</i>	Sea cucumber						
<i>Cucumaria pseudocurata</i>	Sea cucumber						
<i>Dermasterias imbricata</i>	Leather star						
<i>Henricia leviuscula</i>	Blood star						
<i>Leptasterias aequalis</i>							
<i>Leptasterias hexactis</i>	6-rayed star						
<i>Leptasterias puscilla</i>							
<i>Ophiopholis aculeata</i>							
<i>Ophioplocus papillosa</i>							
<i>Ophiothrix spiculata</i>	Brittle star						
<i>Parastichopus parvimensis</i>	Sea cucumber						
<i>Patiria miniata</i>	Bat star						
<i>Pisaster giganteus</i>							
<i>Pisaster ochraceus</i>	Ochre star						
<i>Pycnopodia helianthoides</i>	Sunflower star						
<i>Strongylocentrotus droebachiensis</i>							
<i>Strongylocentrotus franciscanus</i>	Red sea urchin						
<i>Strongylocentrotus purpuratus</i>	Purple sea urchin						

---

**Ectoprocta**

---

Scientific Name	Common Name	Sensitivity to oil	More Benefit from Dispersant	More Harm from Dispersant	ESA listing	Breeds locally	Reason
<i>Barentsia benedeni</i>							

---

<i>Bugula californica</i>	Bryozoan
<i>Crisia maxima</i>	
<i>Dendrobeatia laxa</i>	Bryozoan
<i>Dendrobeatia lichenoides</i>	
<i>Eurystomella bilabiata</i>	
<i>Flustrellidra corniculata</i>	Bryozoan
<i>Tricellaria occidentalis</i>	
<i>Tricellaria sp</i>	
<i>Tricellaria ternata</i>	

---

### Mollusca

---

Scientific Name	Common Name	Sensitivity to oil	More Benefit from Dispersant	More Harm from Dispersant	ESA listing	Breeds locally	Reason
<i>Loligo opalescens</i>		Low	yes	no		yes	Low probability of exposure to oil at the surface, important prey base
<i>Acanthina spirata</i>	Angular unicorn						
<i>Acanthina spp.</i>							
<i>Acanthodoris nanaimoensis</i>							
<i>Aclis shepardiana</i>							
<i>Acmaea mitra</i>	White capped limpet						
<i>Aeolidia papillosa</i>	Shag-rug nudibranch						
<i>Aeolidia papillosa</i>	Nudibranch						
<i>Alia carinata</i>							
<i>Amphissa versicolor</i>	Variegated amphissa						
<i>Anisodoris noblis</i>	Sea lemon						
<i>Antiopella barbarensis</i>							
<i>Archidoris montereyensis</i>	Monterey dorid						
<i>Balcis thersites</i>							
<i>Baptodoris mimetica</i>							
<i>Barleeia haliotiphila</i>	Snail						
<i>Barleeia subtenuis</i>	Snail						
<i>Batillaria attramentaria</i>	Horn snail						
<i>Bittium eschrichtii</i>	Threaded bittium						

<i>Bittium purpureum</i>							
<i>Bittium schrichtii</i>							
<i>Cadlina luteomarginata</i>							
<i>Cadlina modesta</i>	Yellow-edged cadlina						
<i>Calliostoma canaliculatum</i>	Channeled top snail						
<i>Callistoma ligatum</i>	Blue top snail						
<i>Ceratostoma foliatum</i>							
<i>Cerithiopsis carpenteri</i>							
<i>Chama arcana</i>							
<i>Collisella scabra</i>							
<i>Corolla spectabilis (Pteropod)</i>							
<i>Crassostrea gigas</i>	Pacific oyster						
<i>Crepidula adunca</i>	Hooked slipper snail						
<i>Crepidula nummaria</i>							
<i>Crepidula perforans</i>							
<i>Crepidatella lingulata</i>							
<i>Cryptochiton stelleri</i>	Gumboot chiton						
<i>Cryptomya californica</i>							
<i>Cymakra aspera</i>							
<i>Daphana californica</i>							
<i>Diaphana californica</i>							
<i>Diaulula sandiegensis</i>	Ring spotted dorid						
<i>Diplodonta orbella</i>							
<i>Discurria scutum</i>							
<i>Doto columbiana</i>							
<i>Entodesma saxicola</i>							
<i>Epitonium tinctum</i>	Snail						
<i>Fissurella volcano</i>							
<i>Fusinus luteopictus</i>							
<i>Granula margaritula</i>							
<i>Haliotis cracherodii</i>	Black Abalone	High	yes	no	yes	Rare intertidal species likely to be affected by toxicity of oil	
<i>Haliotis rufescens</i>	Red Abalone	High	yes	no	yes	Rare intertidal species likely to be affected by toxicity of oil	

<i>Hermisenda crassicornis</i>	Hermisenda
<i>Hiatella arctica</i>	
<i>Hinnites giganteus</i>	
<i>Hipponix craniodes</i>	Hoof snail
<i>Hopkinsia rosacea</i>	Hopkin's Rose
<i>Irus lamellifer</i>	
<i>Ischnochiton regularis</i>	Chiton
<i>Katharina tunicata</i>	Chiton
<i>Kellia laperousii</i>	
<i>Lacuna cistula</i>	
<i>Lacuna marmorata</i>	Chink snail
<i>Lacuna porrecta</i>	
<i>Lacuna unifasciata</i>	
<i>Lasaea cistula</i>	
<i>Lasaea subviridis</i>	Clam
<i>Lepidochitona dentiens</i>	Chiton
<i>Lepidozona sinudentata</i>	
<i>Littorina keanae</i>	
<i>Littorina planaxis</i>	Eroded periwinkle
<i>Littorina scutulata</i>	Checkered periwinkle
<i>Littorina sitkana</i>	
<i>Littorina sp.</i>	
<i>Lottia asmi</i>	
<i>Lottia digitalis</i>	Ribbed limpet
<i>Lottia gigantea</i>	Owl limpet
<i>Lottia instabilis</i>	Unstable seaweed limpet
<i>Lottia limantula</i>	File limpet
<i>Lottia pelta</i>	Shield limpet
<i>Lottia strigatella</i>	
<i>Lottia triangularis</i>	Triangular limpet
<i>Macclintockia scabra</i>	Rough limpet
<i>Milneria minima</i>	
<i>Mitrella carinata</i>	

<i>Mitrella tuberosa</i>							
<i>Modiolus capax</i>	Fat horse mussel						
<i>Modiolus carpenti</i>							
<i>Mopalia ciliata</i>	Hairy chiton						
<i>Mopalia muscosa</i>	Mossy chiton						
<i>Musculus pygmaeus</i>	Pygmy mussel						
<i>Mytilimeria nuttallii</i>							
<i>Mytilus californianus</i>	California mussel	High	yes	no	yes	Intertidal species likely to be affected by toxicity of oil	
<i>Mytilus edulis</i>	Bay mussel	High	yes	no	yes	Intertidal species likely to be affected by toxicity of oil	
<i>Nassarius mendicus</i>							
<i>Notoacmea insessa</i>	Limpet						
<i>Notoacmea persona</i>	Limpet						
<i>Nucella canaliculata</i>	Channeled dogwinkle						
<i>Nucella emarginata</i>	Emarginate dogwinkle						
<i>Nuttallina californica</i>	Chiton						
<i>Ocenebra atropurpurea</i>							
<i>Ocenebra interfossa</i>							
<i>Ocenebra lurida</i>							
<i>Octopus dofleini</i>							
<i>Octopus rubescens</i>							
<i>Octopus sp.</i>							
<i>Odostomia sp.</i>							
<i>Onchidella borealis</i>							
<i>Opalia wroblewskyi</i>							
<i>Ostrea lurida</i>	Olympic oyster						
<i>Palciphorella velatta</i>							
<i>Penitella conradi</i>							
<i>Penitella turnerae</i>							
<i>Petalochonchus montereyensis</i>							
<i>Petricola carditoides</i>							
<i>Philobrya setosa</i>							
<i>Pododesmus cepio</i>	Abalone jingle						

<i>Protothaca staminea</i>							
<i>Rostanga pulchra</i>	Red sponge nudibranch						
<i>Searlesia dira</i>	Dire welk						
<i>Stenoplax heathiana</i>							
<i>Stiliger fuscovittatus</i>	Streaked stiliger						
<i>Tresus capax</i>		High	yes	no		yes	Intertidal species likely to be affected by toxicity of oil
<i>Saxidomus giganteus</i>		High	yes	no		yes	Intertidal species likely to be affected by toxicity of oil
<i>Siliqua patula</i>		High	yes	no		yes	Intertidal species likely to be affected by toxicity of oil
<i>Tectura insessa</i>							
<i>Tectura persona</i>							
<i>Tectura scutum</i>							
<i>Tegula brunnea</i>	Brown turban snail						
<i>Tegula funebris</i>	Black turban snail						
<i>Tonicella lineata</i>	Lined chiton						
<i>Transennella tantilla</i>							
<i>Trimusculus reticulatus</i>	Reticulate button snail						
<i>Triopha catalinae</i>	Sea-clown nudibranch						
<i>Triopha maculata</i>							
<i>Trivia californica</i>							
<i>Velutina velutina</i>							

---

#### Nemertea

Scientific Name	Common Name	Sensitivity to oil	More Benefit from Dispersant	More Harm from Dispersant	ESA listing	Breeds locally	Reason
<i>Emplectonema gracile</i>							
<i>Tubulanus sexlineatus</i>							

---

#### Porifera

Scientific Name	Common Name	Sensitivity to oil	More Benefit from Dispersant	More Harm from Dispersant	ESA listing	Breeds locally	Reason
<i>Acarus erithacus</i>	Sponge						
<i>Allopora porphyra</i>							
<i>Anaata spongigartina</i>	Sponge						
<i>Antho lithophoenix</i>							
<i>Aplysilla glacialis</i>	Keratose sponge						
<i>Aplysilla polyraphis</i>							
<i>Axocelita originalis</i>	Sponge						
<i>Clathria sp.</i>							
<i>Cliona celata</i>							
<i>Geodia mesotriaence</i>	Sponge						
<i>Halichondria panicea</i>	Crumb-of-bread sponge						
<i>Halichondria sp.</i>							
<i>Haliclona permollis</i>							
<i>Haliclona sp.</i>	Sponge						
<i>Higginsia sp.</i>							
<i>Hinksia sandriana</i>							
<i>Hymedesmia sp.</i>							
<i>Hymenamphiastra cyanocrypta</i>							
<i>Leucandra heathi</i>	Sponge						
<i>Leucilla nuttingi</i>	Sponge						
<i>Leucosolenia eleanor</i>	Sponge						
<i>Lissodendoryx firma</i>	Sponge						
<i>Lissodendoryx topsenti</i>	Sponge						
<i>Mycale psila</i>	Sponge						
<i>Myxilla incrustans</i>							
<i>Ophlitaspongia pennata</i>	Sponge						
<i>Scypha sp.</i>							
<i>Spongia idia</i>							
<i>Stelletta clarella</i>	Sponge						
<i>Suberites sp.</i>	Sponge						

<i>Tedania gurjanovae</i>	Sponge
<i>Tethya aurantia</i>	Sponge
<i>Toxidocia sp.</i>	Sponge
<i>Xestospongia vanilla</i>	Sponge
<i>Zygherpe hyaloderma</i>	Sponge

---

### Sipuncula

Scientific Name	Common Name	Sensitivity to oil	More Benefit from Dispersant	More Harm from Dispersant	ESA listing	Breeds locally	Reason
-----------------	-------------	--------------------	------------------------------	---------------------------	-------------	----------------	--------

*Phascolosoma agassizii*

---

### Urochordata

Scientific Name	Common Name	Sensitivity to oil	More Benefit from Dispersant	More Harm from Dispersant	ESA listing	Breeds locally	Reason
-----------------	-------------	--------------------	------------------------------	---------------------------	-------------	----------------	--------

*Archidistoma ritteri*

*Styela montereyensis*

*Styela truncata*

### PLANTS

---

#### COLOROPHYTA

Scientific Name	Common Name	Sensitivity to oil	More Benefit from Dispersant	More Harm from Dispersant	ESA listing	Breeds locally	Reason
-----------------	-------------	--------------------	------------------------------	---------------------------	-------------	----------------	--------

*Acrosiphonia coalita*

*Bryopsis corticulans* Moss-like algae

*Cladophora columbiana* Pin cushion algae

*Cladophora graminea*

*Cladophora sp.*

*Codium fragile* Dead man's fingers

<i>Codium setchellii</i>	Sponge weed
<i>Derbesia marina</i>	
<i>Endocladia viridis</i>	
<i>Endophyton ramosum</i>	
<i>Enteromorpha flexuosa</i>	
<i>Enteromorpha clathrata</i>	
<i>Enteromorpha compressa</i>	
<i>Enteromorpha intestinalis</i>	Intestine alge
<i>Halicystis ovalis</i>	
<i>Prasiola meridionalis</i>	
<i>Ulothrix flacca</i>	
<i>Ulothrix laetevirens</i>	
<i>Ulothrix pseudoflacca</i>	
<i>Ulva californica</i>	
<i>Ulva conglobata</i>	
<i>Ulva expansa</i>	
<i>Ulva lactuca</i>	
<i>Ulva lobata</i>	
<i>Ulva spp.</i>	Sea lettuce
<i>Ulva taeniata</i>	
<i>Urophoro sp.</i>	

---

**HETEROKONTOPHYTA**

---

Scientific Name	Common Name	Sensitivity to oil	More Benefit from Dispersant	More Harm from Dispersant	ESA listing	Breeds locally	Reason
<i>Alaria marginata</i>	Winged kelp						
<i>Anelipes japonicus</i>	Barefoot, Matsumo						
<i>Coilodesme californica</i>							
<i>Colpomenia peregrina</i>							
<i>Compsonea serpens</i>							
<i>Costaria costata</i>							
<i>Cystoseira osmundacea</i>	Bladder chain						

<i>Desmarestia herbacea</i>							
<i>Desmarestia ligulata</i>	Acid seaweed						
<i>Desmarestia munda</i>							
<i>Dictyoneurum californicum</i>	Nerve net						
<i>Egregia menziesii</i>	Feather Boa						
<i>Fucus gardneri</i>	Rock weed	High	Yes	No			Experience of injury documented in intertidal monitoring plots
<i>Hinckesia sandriana</i>							
<i>Laminaria ephemera</i>							
<i>Laminaria farlowii</i>							
<i>Laminaria setchellii</i>	Split blade oarweed/Kombu						
<i>Laminaria sinclarii</i>	Oar weed/Kombu						
<i>Laminaria sp.</i>							
<i>Leathesia difformis</i>							
<i>Macrocystis integrifolia</i>							
<i>Macrocystis pyrifera</i>	Giant Kelp	Medium	Yes	No	yes		Hold fast protected at depth, mucus protects kelp, kelp may retain oil and increase toxic exposure of organisms in that habitat
<i>Melanosiphon intestinalis</i>							
<i>Nereocystis luetkeana</i>	Bull whip kelp						
<i>Nereocystis luetkeana</i>	Bull Kelp	Medium	Yes	No	yes		Hold fast protected at depth, mucus protects kelp, kelp may retain oil and increase toxic exposure of organisms in that habitat
<i>Pelvetia fastigiata</i>	Little rock weed						
<i>Pelvetiopsis limitata</i>	Tiny rock weed						
<i>Petalonia fascia</i>							
<i>Phaeostrophion irregulare</i>							
<i>Pilayella sp.</i>							
<i>Postelsia palmaeformis</i>	Sea palm	High	Yes	No	yes		Hold fast exposed at intertidal, kelp may retain oil and increase toxic exposure of organisms in that habitat
<i>Pterygophora californica</i>							
<i>Ralfsia pacifica</i>	Tar spot						

*Ralfsia sp.*  
*Sargassum muticum*  
*Scytisiphon simplicissimus*      Leather tube  
*Scytosiphon dotyii*  
*Scytosiphon lomentaria*  
*Soranothera ulvoidea*  
*Spongonema tomentosum*  
*Streblonema sp.*

---

**RHODOPHYTA**

---

Scientific Name	Common Name	Sensitivity to oil	More Benefit from Dispersant	More Harm from Dispersant	ESA listing	Breeds locally	Reason
<i>Acrochaetium prophyrae</i>	Dreadlock algae						
<i>Acrochaetium sp.</i>	Epiphytic algae						
<i>Ahnfeltia cornucopiae</i>	Garlic algae						
<i>Ahnfeltia fastigiata</i>	Mastocarpus crust						
<i>Ahnfeltiopsis leptophylla</i>							
<i>Ahnfeltiopsis linearis</i>							
<i>Anotrichium furcellatum</i>	Red membrane						
<i>Antithamnion dendroidum</i>							
<i>Antithamnion densum</i>							
<i>Audouinella subimmersa</i>	Tooth branch						
<i>Bangia sp.</i>	Braided hair algae						
<i>Bornetia californica</i>							
<i>Bossiella corymbifera</i>							
<i>Bossiella dichotoma</i>							
<i>Bossiella plumosa</i>							
<i>Bossiella schmittii</i>							
<i>Branchioglossum bipinnatifidum</i>							
<i>Branchioglossum undulatum</i>							
<i>Callithamnion biseriatum</i>							
<i>Callophyllis cheilosporioides</i>							

*Callophyllis crenulata*  
*Callophyllis flabellulata*  
*Callophyllis heanophylla*  
*Callophyllis linearis*  
*Callophyllis obtusifolia*  
*Callophyllis pinnata*  
*Callophyllis sp.*  
*Callophyllis violacea*  
*Centroceras clavulatum*  
*Ceramium gardneri*  
*Ceramium pacificum*  
*Chiharaea bodegensis*  
*Cirriliarpus sp.*  
*Clathromorphum parcum*  
*Constantinea simplex*  
*Corallina officinalis*  
*Corallina pinnatifolia*  
*Crustose corallines*  
*Cryptopleura farlowiana*  
*Cryptopleura corallinara*  
*Cryptopleura crispa*  
*Cryptopleura lobulifera*  
*Cryptopleura rosacea*  
*Cryptopleura ruprechtiana*  
*Cumagloia andersonii*  
*Delesseria decipiens*  
*Dilsea californica*  
*Endocladia muricata*                      Beautifully jointed  
*Erythrogloum californicum*  
*Erythrophyllum delesseriodes*              Wool weed  
*Erythrotrichia carnea*  
*Erythrotrichia pulvinata*  
*Farlowia compressa*

<i>Farlowia conferta</i>	
<i>Farlowia mollis</i>	
<i>Faucheia fryeana</i>	
<i>Faucheia laciniata</i>	
<i>Faucheocolax attenuata</i>	
<i>Gastroclonium subarticulatum</i>	Beautiful leaf
<i>Gastroclonium subarticulatum</i>	
<i>Gelidium coulteri</i>	Candy cane seaweed
<i>Gelidium purpurascens</i>	Arrow weed
<i>Gelidium pusillum</i>	
<i>Gelidium robustum</i>	
<i>Gelidium sp.</i>	
<i>Gloiosiphonia verticillaris</i>	
<i>Goniotrichopsis sublittoralis</i>	
<i>Gracilariophila oryzoides</i>	
<i>Gracilariopsis sjoestedtii</i>	Turkish towel
<i>Grateloupia doryphora</i>	
<i>Grateloupia filicina</i>	
<i>Griffithsia pacifica</i>	
<i>Gymnogongrus chiton</i>	
<i>Halosaccion glandiforme</i>	Turkish towel
<i>Halymenia schizymenioides</i>	
<i>Halymenia templetonii</i>	
<i>Herposiphonia parva</i>	
<i>Herposiphonia plumula</i>	
<i>Hildenbrandia occidentalis</i>	
<i>Hildenbrandia rubra</i>	
<i>Hildenbrandia spp.</i>	Narrow turkish towel
<i>Hommersandia palmatifolia</i>	
<i>Hymenena coccinea</i>	
<i>Hymenena flabelligera</i>	
<i>Hymenena multiloba</i>	
<i>Janczewskia gardneri</i>	

<i>Leachiella pacifica</i>	
<i>Lithophyllum dispar</i>	
<i>Lithophyllum grumosum</i>	
<i>Lithophyllum proboscideum</i>	
<i>Lithothamnium sp.</i>	Narrow turkish towel
<i>Lithothrix aspergillum</i>	Cup and saucer algae
<i>Maripelta rotata</i>	
<i>Mastocarpus jardinii</i>	Small coral
<i>Mastocarpus papillatus</i>	Hidden ribs
<i>Mazzaella affinis</i>	
<i>Mazzaella californica</i>	
<i>Mazzaella cordata</i>	
<i>Mazzaella cornucopiae</i>	Nail brush
<i>Mazzaella flaccida</i>	Red leaf
<i>Mazzaella heterocarpa</i>	Belly branch
<i>Mazzaella leptorhynchos</i>	
<i>Mazzaella linearis</i>	
<i>Mazzaella rosea</i>	
<i>Mazzaella splendens</i>	Agarweed
<i>Mazzaella volans</i>	
<i>Melobesia marginata</i>	
<i>Melobesia mediocris</i>	Agarweed
<i>Membranoptera dimorpha</i>	
<i>Mesophyllum conchatum</i>	
<i>Mesophyllum lamellatum</i>	
<i>Microcladia borealis</i>	Spaghetti weed
<i>Microcladia coulteri</i>	Sea sac
<i>Myriogramme sp.</i>	
<i>Myriogramme spectabilis</i>	
<i>Myriogramme variegata</i>	
<i>Neoptilota densa</i>	
<i>Neoptilota hypnoides</i>	
<i>Neoptilota sp.</i>	

<i>Neorhodomela larix</i>	Wine crust
<i>Nienburgia andersoniana</i>	
<i>Nitophyllum sp.</i>	
<i>Nitophyllum sp.</i>	
<i>Odonthalia floccosa</i>	crustose coralline
<i>Opuntiella californica</i>	Stone hair
<i>Osmundea spectabilis</i>	Little turkish towel
<i>Petrocelis franciscana</i>	Little turkish towel
<i>Petrospongium rugosum</i>	
<i>Peyssonelliopsis epiphytica</i>	
<i>Peyssonnelia meridionalis</i>	
<i>Peyssonnelia pacifica</i>	
<i>Phycodryes setchellii</i>	
<i>Pikea californica</i>	
<i>Pikea pinnata</i>	
<i>Pleonosporium</i>	
<i>vancouverianum</i>	
<i>Plocamium cartilagineum</i>	Bunny ears algae
<i>Plocamium cartilagineum var. pacificum</i>	
<i>Plocamium oregonum</i>	
<i>Plocamium pacificum</i>	
<i>Plocamium sp.</i>	
<i>Plocamium violaceum</i>	
<i>Polyneura latissima</i>	Iridesent seaweed
<i>Polysiphonia hendryi</i>	Warty algae
<i>Polysiphonia hendryi</i>	
<i>Polysiphonia pacifica</i>	
<i>Polysiphonia saraticeri</i>	
<i>Polysiphonia sp.</i>	
<i>Porphyra gardneri</i>	Many veined algae
<i>Porphyra lanceolata</i>	Many siphon algae
<i>Porphyra nereocystis</i>	Nori/laver
<i>Porphyra perforata</i>	Iridesent seaweed
<i>Porphyra sp.</i>	Serrated red weed

<i>Prionitis australis</i>	
<i>Prionitis cornea</i>	
<i>Prionitis lanceolata</i>	Phyllospadix crust
<i>Prionitis linearis</i>	
<i>Prionitis lyallii</i>	
<i>Pronitis filiformis</i>	
<i>Pronitis sp.</i>	
<i>Pseudolithophyllum neofarlowii</i>	
<i>Pterochondria woodii</i>	
<i>Pteroclatiella caloglossoides</i>	
<i>Pteroclatiella capillacea</i>	
<i>Pterosiphonia baileyi</i>	
<i>Pterosiphonia bipinnata</i>	
<i>Pterosiphonia dendroidea</i>	
<i>Pterothamnion villosum</i>	
<i>Ptilota filicina</i>	
<i>Ptilothamnionopsis lejolisea</i>	
<i>Pugetia fragilissima</i>	
<i>Rhodochorton purpureum</i>	Cactus weed
<i>Rhodymenia californica</i>	Small branch
<i>Rhodymenia callophyllidoides</i>	
<i>Rhodymenia pacifica</i>	
<i>Sahlingia subintegra</i>	
<i>Sarcodiotheca gaudichaudii</i>	
<i>Schimmelemannia plumosa</i>	
<i>Schizymenia pacifica</i>	
<i>Scinaia confusa</i>	
<i>Smithora naiadum</i>	
<i>Stenogramma interrupta</i>	
<i>Stylonema alsidii</i>	
<i>Tiffaniella snyderae</i>	
<i>Titanoderma dispar</i>	
<i>Weeksia reticulata</i>	

---

**VASCULAR**

---

Scientific Name	Common Name	Sensitivity to oil	More Benefit from Dispersant	More Harm from Dispersant	ESA listing	Breeds locally	Reason
<i>Phyllospadix scouleri</i>	Surf grass	Medium	Yes	No		yes	Grass may retain oil and increase toxic exposure of organisms in that habitat
<i>Phyllospadix torreyi</i>		Medium	Yes	No		yes	Grass may retain oil and increase toxic exposure of organisms in that habitat
<i>Zostera marina</i>	Eel grass	Medium	Yes	No		yes	Grass may retain oil and increase toxic exposure of organisms in that habitat

## Appendix V: Sensitive Species Matrix

Feb15-Aug15				Cordell Bank	Farallon Scarpment (a.k.a FNWR from Fanny Shoals to SEFI)	Deep Reef	Point Reyes	Drakes Bay	Devil's Slide	PR Convergent Zone	SF Bay Plume	Dunes and Beaches	Open Estuaries (SF, Tomales, Bolinas)	Seasonal Estuaries (Rodeo, San Antonio etc)	Tidepools (Duxbury, Fitzgerald, etc)	
	COMMON NAME	SCIENTIFIC NAME	Benefit from dispersants?													
Mammals	Gray Whale	<i>Eschrichtius robustus</i>		-	1	-	1	1	1	1	1	-	1	-	-	-
Mammals	Humpback Whale	<i>Megaptera novaeangliae</i>		1	1	1	1	1	1	1	1	-	-	-	-	-
Mammals	Sea Otter	<i>Enhydra lutris</i>		-	-	-	-	-	1	-	1	-	-	-	-	1
Mammals	Harbor Porpoise	<i>Phocoena phocoena</i>		-	-	-	1	1	1	1	1	-	1	-	-	-
Mammals	Northern Fur Seal	<i>Callorhinus ursinus</i>		1	1	1	-	-	-	1	-	-	-	-	-	-
Mammals	Harbor Seal	<i>Phoca vitulina</i>		-	1	-	1	1	1	1	1	1	1	-	-	1
Bird	Ashy Storm-Petrel	<i>Oceanodroma homochroa</i>		1	1	1	1	1	-	1	1	-	-	-	-	-
Bird	Cassin's Auklet	<i>Ptychoramphus aleuticus</i>		1	1	1	-	-	-	1	1	-	-	-	-	-
Bird	Snowy Plover	<i>Charadrius alexandrinus</i>		-	-	-	-	-	-	-	-	1	-	-	-	-
Bird	Marbled Murrelet,	<i>Brachyramphus marmoratus</i>		-	-	-	1	1	1	1	1	-	-	-	-	-
Bird	Sooty Shearwater	<i>Puffinus griseus</i>		1	1	1	-	-	-	1	1	-	-	-	-	-
Bird	Surf Scoter	<i>Melanitta perspicillata</i>		-	-	-	1	1	1	1	1	-	1	-	-	-
Bird	Brandt's Cormorant	<i>Phalacrocorax penicillatus</i>		-	1	-	1	1	1	1	1	-	1	-	-	1
Bird	Common Murre	<i>Uria aalge</i>		1	1	1	1	1	1	1	1	-	1	-	-	-
Bird	Osprey	<i>Pandion haliaetus</i>		-	-	-	1	1	-	-	1	1	-	1	-	-
Bird	Peregrine	<i>Falco peregrinus</i>		-	1	-	1	1	1	-	-	1	1	1	1	1
Fish	Tidewater Goby	<i>Eucyclogobius newberryi</i>		-	-	-	-	-	1	-	-	-	1	1	1	1
Fish	Coho Salmon [Silver Salmon]	<i>Oncorhynchus kisutch</i>		1	1	1	1	1	1	1	1	-	1	1	1	-
Fish	Pacific Herring	<i>Clupea pallasii</i>		-	-	-	1	1	-	-	-	-	1	-	-	-
Fish	Rockfish (Juveniles)			1	1	1	1	1	1	1	1	-	-	-	-	-
Crustacea	Krill	<i>Euphausia pacifica</i>		1	1	1	1	-	-	-	-	-	-	-	-	-
Crustacea	Dungeness crab	<i>Cancer magister</i>		-	-	1	1	1	1	-	-	-	1	1	-	-
Mollusca	Red Abalone	<i>Haliotis rufescens</i>		-	1	-	1	1	1	-	-	-	-	-	-	1
Mollusca	California mussel	<i>Mytilus californianus</i>		-	1	-	1	1	1	-	-	-	1	-	-	1
Algae	Rock weed	<i>Fucus gardneri</i>		-	1	-	1	-	1	-	-	-	-	-	-	1
Algae	Sea palm	<i>Postelsia palmaeformis</i>		-	1	-	1	-	1	-	-	-	-	-	-	1
Vascular	Surf grass	<i>Phyllospadix scouleri</i>		-	-	-	-	1	-	-	-	-	1	-	-	1
Vascular	Eel grass	<i>Zostera marina</i>		-	-	-	-	1	-	-	-	-	1	-	-	-

# Aug15-Nov15

Aug15-Nov15				Cordell Bank	Farallon Scarpment (a.k.a FNWR from Fanny Shoals to SEFI)	Deep Reef	Point Reyes	Drakes Bay	Devil's Slide	PR Convergent Zone	SF Bay Plume	Dunes and Beaches	Open Estuaries (SF, Tomales, Bolinas)	Seasonal Estuaries (Rodeo, San Antonio etc)	Tidepools (Duxbury, Fitzgerald, etc)	
	COMMON NAME	SCIENTIFIC NAME	Benefit from dispersants?													
Mammals	Gray Whale	<i>Eschrichtius robustus</i>		-	1	-	1	1	1	1	1	-	1	-	-	
Mammals	Humpback Whale	<i>Megaptera novaeangliae</i>		1	1	1	1	1	1	1	1	-	-	-	-	
Mammals	Sea Otter	<i>Enhydra lutris</i>		-	-	-	-	-	1	-	1	-	-	-	1	
Mammals	Harbor Porpoise	<i>Phocoena phocoena</i>		-	-	-	1	1	1	1	1	-	1	-	-	
Mammals	Northern Fur Seal	<i>Callorhinus ursinus</i>		1	1	1	-	-	-	1	-	-	-	-	-	
Mammals	Harbor Seal	<i>Phoca vitulina</i>		-	1	-	1	1	1	1	1	1	1	-	1	
Bird	Ashy Storm-Petrel	<i>Oceanodroma homochroa</i>		-	-	-	-	-	1	-	-	-	1	1	1	
Bird	Cassin's Auklet	<i>Ptychoramphus aleuticus</i>		1	1	1	1	1	1	1	1	-	1	1	-	
Bird	Snowy Plover	<i>Charadrius alexandrinus</i>		-	-	-	1	1	-	-	-	-	1	-	-	
Bird	Marbled Murrelet	<i>Brachyramphus marmoratus</i>		1	1	1	1	1	1	1	1	-	-	-	-	
Bird	Sooty Shearwater	<i>Puffinus griseus</i>		1	1	1	1	-	-	-	-	-	-	-	-	
Bird	Surf Scoter	<i>Melanitta perspicillata</i>		-	-	1	1	1	1	-	-	-	1	1	-	
Bird	Brandt's Cormorant	<i>Phalacrocorax penicillatus</i>		-	1	-	1	1	1	-	-	-	-	-	1	
Bird	Common Murre	<i>Uria aalge</i>		-	1	-	1	1	1	-	-	-	1	-	1	
Bird	Osprey	<i>Pandion haliaetus</i>		-	-	-	1	1	-	-	1	1	1	1	-	
Bird	Peregrine	<i>Falco peregrinus</i>		-	1	-	1	1	1	-	-	1	1	1	1	
Fish	Tidewater Goby	<i>Eucyclogobius newberryi</i>		-	1	-	1	-	1	-	-	-	-	-	-	1
Fish	Coho Salmon [Silver Salmon]	<i>Oncorhynchus kisutch</i>		-	1	-	1	-	1	-	-	-	-	-	-	1
Fish	Pacific Herring	<i>Clupea pallasii</i>		-	-	-	-	1	-	-	-	-	1	-	1	
Fish	Juvenile Rockfish			-	-	-	-	1	-	-	-	-	1	-	-	
Crustacea	Krill	<i>Euphausia pacifica</i>		1	1	1	1	-	-	-	-	-	-	-	-	
Crustacea	Dungeness crab	<i>Cancer magister</i>		-	-	1	1	1	1	-	-	-	1	1	-	
Mollusca	Red Abalone	<i>Haliotis rufescens</i>		-	1	-	1	1	1	-	-	-	-	-	-	1
Mollusca	California mussel	<i>Mytilus californianus</i>		-	1	-	1	1	1	-	-	-	1	-	1	
Algae	Rock weed	<i>Fucus gardneri</i>		-	1	-	1	-	1	-	-	-	-	-	-	1
Algae	Sea palm	<i>Postelsia palmaeformis</i>		-	1	-	1	-	1	-	-	-	-	-	-	1
Vascular	Surf grass	<i>Phyllospadix scouleri</i>		-	-	-	-	1	-	-	-	-	1	-	1	
Vascular	Eel grass	<i>Zostera marina</i>		-	-	-	-	1	-	-	-	-	1	-	-	

# Nov15-Feb15

Nov15-Feb15				Cordell Bank	Farallon Scarpment (a.k.a FNWR from Fanny Shoals to SEFI)	Deep Reef	Point Reyes	Drakes Bay	Devil's Slide	PR Convergent Zone	SF Bay Plume	Dunes and Beaches	Open Estuaries (SF, Tomales, Bolinas)	Seasonal Estuaries (Rodeo, San Antonio etc)	Tidepools (Duxbury, Fitzgerald, etc)	
	COMMON NAME	SCIENTIFIC NAME	Benefit from dispersants?													
Mammals	Gray Whale	<i>Eschrichtius robustus</i>		-	1	-	1	1	1	1	1	-	1	-	-	
Mammals	Humpback Whale	<i>Megaptera novaeangliae</i>		1	1	1	1	1	1	1	1	-	-	-	-	
Mammals	Sea Otter	<i>Enhydra lutris</i>		-	-	-	-	-	1	-	1	-	-	-	1	
Mammals	Harbor Porpoise	<i>Phocoena phocoena</i>		-	-	-	1	1	1	1	1	-	1	-	-	
Mammals	Northern Fur Seal	<i>Callorhinus ursinus</i>		1	1	1	-	-	-	1	-	-	-	-	-	
Mammals	Harbor Seal	<i>Phoca vitulina</i>		-	1	-	1	1	1	1	1	1	1	-	1	
Bird	Ashy Storm-Petrel	<i>Oceanodroma homochroa</i>		-	-	-	-	-	1	-	-	-	1	1	1	
Bird	Cassin's Auklet	<i>Ptychoramphus aleuticus</i>		1	1	1	1	1	1	1	1	-	1	1	-	
Bird	Snowy Plover	<i>Charadrius alexandrinus</i>		-	-	-	1	1	-	-	-	-	1	-	-	
Bird	Marbled Murrelet,	<i>Brachyramphus marmoratus</i>		1	1	1	1	1	1	1	1	-	-	-	-	
Bird	Sooty Shearwater	<i>Puffinus griseus</i>		1	1	1	1	-	-	-	-	-	-	-	-	
Bird	Surf Scoter	<i>Melanitta perspicillata</i>		-	-	1	1	1	1	-	-	-	1	1	-	
Bird	Brandt's Cormorant	<i>Phalacrocorax penicillatus</i>		-	1	-	1	1	1	-	-	-	-	-	-	1
Bird	Common Murre	<i>Uria aalge</i>		-	1	-	1	1	1	-	-	-	1	-	1	
Bird	Osprey	<i>Pandion haliaetus</i>		-	-	-	1	1	-	-	1	1	1	1	-	
Bird	Peregrine	<i>Falco peregrinus</i>		-	1	-	1	1	1	-	-	1	1	1	1	
Fish	Tidewater Goby	<i>Eucyclogobius newberryi</i>		-	1	-	1	-	1	-	-	-	-	1	1	
Fish	Coho Salmon [Silver Salmon]	<i>Oncorhynchus kisutch</i>		-	1	-	1	-	1	-	-	-	1	1	1	
Fish	Pacific Herring	<i>Clupea pallasii</i>		-	-	-	-	1	-	-	-	-	1	-	1	
Fish	Juvenile Rockfish			-	-	-	-	1	-	-	-	-	1	-	-	
Crustacea	Krill	<i>Euphausia pacifica</i>		1	1	1	1	-	-	-	-	-	-	-	-	
Crustacea	Dungeness crab	<i>Cancer magister</i>		-	-	1	1	1	1	-	-	-	1	1	-	
Mollusca	Red Abalone	<i>Haliotis rufescens</i>		-	1	-	1	1	1	-	-	-	-	-	1	
Mollusca	California mussel	<i>Mytilus californianus</i>		-	1	-	1	1	1	-	-	-	1	-	1	
Algae	Rock weed	<i>Fucus gardneri</i>		-	1	-	1	-	1	-	-	-	-	-	-	1
Algae	Sea palm	<i>Postelsia palmaeformis</i>		-	1	-	1	-	1	-	-	-	-	-	-	1
Vascular	Surf grass	<i>Phyllospadix scouleri</i>		-	-	-	-	1	-	-	-	-	1	-	1	
Vascular	Eel grass	<i>Zostera marina</i>		-	-	-	-	1	-	-	-	-	1	-	1	